H Answers to Odd-Numbered Exercises

CHAPTER 10

EXERCISES 10.1 ■ PAGE 665

1.



3.



5. (a)



(b) $y = \frac{3}{4}x - \frac{1}{4}$

7. (a)



(b) $x = -(y + 2)^2 + 1$, $-4 \leqslant y \leqslant 0$

9. (a)



(b) $y = 1 - x^2$, $x \geqslant 0$

11. (a) $x^2 + y^2 = 1$, $y \geqslant 0$ (b)



13. (a) $y = 1/x$, $y > 1$ (b)



15. (a) $y = \frac{1}{2}\ln x + 1$ (b)



17. (a) $y^2 - x^2 = 1$, $y \geqslant 1$

(b)



19. Moves counterclockwise along the circle $(x - 3)^2 + (y - 1)^2 = 4$ from (3, 3) to (3, −1)

21. Moves 3 times clockwise around the ellipse $(x^2/25) + (y^2/4) = 1$, starting and ending at (0, −2)

23. It is contained in the rectangle described by $1 \leqslant x \leqslant 4$ and $2 \leqslant y \leqslant 3$.

25.



27.



29.



31. (b) $x = -2 + 5t$, $y = 7 - 8t$, $0 \leqslant t \leqslant 1$

33. (a) $x = 2\cos t$, $y = 1 - 2\sin t$, $0 \leqslant t \leqslant 2\pi$

(b) $x = 2\cos t$, $y = 1 + 2\sin t$, $0 \leqslant t \leqslant 6\pi$

(c) $x = 2\cos t$, $y = 1 + 2\sin t$, $\pi/2 \leqslant t \leqslant 3\pi/2$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

37. The curve $y = x^{2/3}$ is generated in (a). In (b), only the portion with $x \geqslant 0$ is generated, and in (c) we get only the portion with $x > 0$.

41. $x = a\cos\theta, y = b\sin\theta$; $(x^2/a^2) + (y^2/b^2) = 1$, ellipse

43.



45. (a) Two points of intersection



(b) One collision point at $(-3, 0)$ when $t = 3\pi/2$

(c) There are still two intersection points, but no collision point.

47. For $c = 0$, there is a cusp; for $c > 0$, there is a loop whose size increases as c increases.





49. The curves roughly follow the line $y = x$, and they start having loops when a is between 1.4 and 1.6 (more precisely, when $a > \sqrt{2}$). The loops increase in size as a increases.

51. As n increases, the number of oscillations increases; a and b determine the width and height.

EXERCISES 10.2 ■ PAGE 675

1. $\dfrac{2t + 1}{t\cos t + \sin t}$ **3.** $y = -\frac{3}{2}x + 7$ **5.** $y = \pi x + \pi^2$

7. $y = 2x + 1$

9. $y = \frac{1}{6}x$



11. $\dfrac{2t + 1}{2t}, -\dfrac{1}{4t^3}, t < 0$ **13.** $e^{-2t}(1 - t), e^{-3t}(2t - 3), t > \frac{3}{2}$

15. $-\frac{3}{2}\tan t, -\frac{3}{4}\sec^3 t, \pi/2 < t < 3\pi/2$

17. Horizontal at $(0, -3)$, vertical at $(\pm 2, -2)$

19. Horizontal at $\left(\frac{1}{2}, -1\right)$ and $\left(-\frac{1}{2}, 1\right)$, no vertical

21. $(0.6, 2)$; $\left(5 \cdot 6^{-6/5}, e^{6^{-1/5}}\right)$

23.



25. $y = x, y = -x$



27. (a) $d\sin\theta/(r - d\cos\theta)$ **29.** $\left(\frac{16}{27}, \frac{29}{9}\right), (-2, -4)$

31. πab **33.** $3 - e$ **35.** $2\pi r^2 + \pi d^2$

37. $\int_0^2 \sqrt{2 + 2e^{-2t}}\, dt \approx 3.1416$

39. $\int_0^{4\pi} \sqrt{5 - 4\cos t}\, dt \approx 26.7298$ **41.** $4\sqrt{2} - 2$

43. $\frac{1}{2}\sqrt{2} + \frac{1}{2}\ln\left(1 + \sqrt{2}\right)$

45. $\sqrt{2}\,(e^{\pi} - 1)$



47. 16.7102



49. 612.3053 **51.** $6\sqrt{2}, \sqrt{2}$

55. (a) $t \in [0, 4\pi]$



(b) 294

57. $\int_0^{\pi/2} 2\pi t\cos t\sqrt{t^2 + 1}\, dt \approx 4.7394$

59. $\int_0^1 2\pi(t^2 + 1)e^t\sqrt{e^{2t}(t + 1)^2(t^2 + 2t + 2)}\, dt \approx 103.5999$

61. $\frac{2}{1215}\pi\left(247\sqrt{13} + 64\right)$ **63.** $\frac{6}{5}\pi a^2$

65. $\frac{24}{5}\pi\left(949\sqrt{26} + 1\right)$ **71.** $\frac{1}{4}$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

EXERCISES 10.3 ■ PAGE 686

1. (a)



(b)



$(2, 7\pi/3), (-2, 4\pi/3)$ $(1, 5\pi/4), (-1, \pi/4)$

(c)



$(1, 3\pi/2), (-1, 5\pi/2)$

3. (a)



$(-1, 0)$

(b)



$\left(-1, -\sqrt{3}\right)$

(c)



$\left(\sqrt{2}, -\sqrt{2}\right)$

5. (a) (i) $\left(2\sqrt{2}, 7\pi/4\right)$ (ii) $\left(-2\sqrt{2}, 3\pi/4\right)$
(b) (i) $(2, 2\pi/3)$ (ii) $(-2, 5\pi/3)$

7.



9.



11.



13. $2\sqrt{3}$ **15.** Circle, center O, radius $\sqrt{5}$
17. Circle, center $(1, 0)$, radius 1
19. Hyperbola, center O, foci on x-axis
21. $r = 2 \csc \theta$ **23.** $r = 1/(\sin \theta - 3 \cos \theta)$
25. $r = 2c \cos \theta$ **27.** (a) $\theta = \pi/6$ (b) $x = 3$

29.



31.



33.



35.



37.



39.



41.



43.



45.



47.



Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

49



51.



53. (a) For $c < -1$, the inner loop begins at $\theta = \sin^{-1}(-1/c)$ and ends at $\theta = \pi - \sin^{-1}(-1/c)$; for $c > 1$, it begins at $\theta = \pi + \sin^{-1}(1/c)$ and ends at $\theta = 2\pi - \sin^{-1}(1/c)$.

55. $\sqrt{3}$ **57.** $-\pi$ **59.** 1

61. Horizontal at $\left(3/\sqrt{2},\ \pi/4\right)$, $\left(-3/\sqrt{2},\ 3\pi/4\right)$; vertical at $(3,\ 0)$, $(0,\ \pi/2)$

63. Horizontal at $\left(\frac{3}{2},\ \pi/3\right)$, $(0,\ \pi)$ [the pole], and $\left(\frac{3}{2},\ 5\pi/3\right)$; vertical at $(2,\ 0)$, $\left(\frac{1}{2},\ 2\pi/3\right)$, $\left(\frac{1}{2},\ 4\pi/3\right)$

65. Center $(b/2,\ a/2)$, radius $\sqrt{a^2 + b^2}/2$

67.



69.



71.



73. By counterclockwise rotation through angle $\pi/6$, $\pi/3$, or α about the origin

75. For $c = 0$, the curve is a circle. As c increases, the left side gets flatter, then has a dimple for $0.5 < c < 1$, a cusp for $c = 1$, and a loop for $c > 1$.

EXERCISES 10.4 ■ PAGE 692

1. $e^{-\pi/4} - e^{-\pi/2}$ **3.** $\frac{9}{2}$ **5.** π^2 **7.** $\frac{41}{4}\pi$

9. π



11. 11π



13. $\frac{9}{2}\pi$



15. $\frac{3}{2}\pi$



17. $\frac{4}{3}\pi$ **19.** $\frac{1}{16}\pi$ **21.** $\pi - \frac{3}{2}\sqrt{3}$ **23.** $\frac{1}{3}\pi + \frac{1}{2}\sqrt{3}$

25. $4\sqrt{3} - \frac{4}{3}\pi$ **27.** π **29.** $\frac{5}{24}\pi - \frac{1}{4}\sqrt{3}$ **31.** $\frac{1}{2}\pi - 1$

33. $1 - \frac{1}{2}\sqrt{2}$ **35.** $\frac{1}{4}\left(\pi + 3\sqrt{3}\right)$

37. $\left(\frac{3}{2},\ \pi/6\right)$, $\left(\frac{3}{2},\ 5\pi/6\right)$, and the pole

39. $(1,\ \theta)$ where $\theta = \pi/12,\ 5\pi/12,\ 13\pi/12,\ 17\pi/12$ and $(-1,\ \theta)$ where $\theta = 7\pi/12,\ 11\pi/12,\ 19\pi/12,\ 23\pi/12$

41. $\left(\frac{1}{2}\sqrt{3},\ \pi/3\right)$, $\left(\frac{1}{2}\sqrt{3},\ 2\pi/3\right)$, and the pole

43. Intersection at $\theta \approx 0.89,\ 2.25$; area ≈ 3.46

45. 2π **47.** $\frac{8}{3}\left[(\pi^2 + 1)^{3/2} - 1\right]$

49. $\frac{16}{3}$



51. 2.4221 **53.** 8.0091

55. (b) $2\pi\left(2 - \sqrt{2}\right)$

EXERCISES 10.5 ■ PAGE 700

1. $(0,\ 0)$, $\left(0,\ \frac{3}{2}\right)$, $y = -\frac{3}{2}$



3. $(0,\ 0)$, $\left(-\frac{1}{2},\ 0\right)$, $x = \frac{1}{2}$



5. $(-2,\ 3)$, $(-2,\ 5)$, $y = 1$



7. $(-2,\ -1)$, $(-5,\ -1)$, $x = 1$



9. $x = -y^2$, focus $\left(-\frac{1}{4},\ 0\right)$, directrix $x = \frac{1}{4}$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

11. $(0, \pm 2), \left(0, \pm\sqrt{2}\right)$



13. $(\pm 3, 0), \left(\pm 2\sqrt{2}, 0\right)$



15. $(1, \pm 3), \left(1, \pm\sqrt{5}\right)$



17. $\dfrac{x^2}{4} + \dfrac{y^2}{9} = 1$, foci $\left(0, \pm\sqrt{5}\right)$

19. $(0, \pm 5); \left(0, \pm\sqrt{34}\right); y = \pm\frac{5}{3}x$



21. $(\pm 10, 0), \left(\pm 10\sqrt{2}, 0\right), y = \pm x$



23. $(4, -2), (2, -2)$; $\left(3 \pm \sqrt{5}, -2\right)$; $y + 2 = \pm 2(x - 3)$



25. Parabola, $(0, -1), \left(0, -\frac{3}{4}\right)$

27. Ellipse, $\left(\pm\sqrt{2}, 1\right), (\pm 1, 1)$

29. Hyperbola, $(0, 1), (0, -3); \left(0, -1 \pm \sqrt{5}\right)$

31. $y^2 = 4x$ **33.** $y^2 = -12(x + 1)$ **35.** $y - 3 = 2(x - 2)^2$

37. $\dfrac{x^2}{25} + \dfrac{y^2}{21} = 1$ **39.** $\dfrac{x^2}{12} + \dfrac{(y-4)^2}{16} = 1$

41. $\dfrac{(x+1)^2}{12} + \dfrac{(y-4)^2}{16} = 1$ **43.** $\dfrac{x^2}{9} - \dfrac{y^2}{16} = 1$

45. $\dfrac{(y-1)^2}{25} - \dfrac{(x+3)^2}{39} = 1$ **47.** $\dfrac{x^2}{9} - \dfrac{y^2}{36} = 1$

49. $\dfrac{x^2}{3{,}763{,}600} + \dfrac{y^2}{3{,}753{,}196} = 1$

51. (a) $\dfrac{121x^2}{1{,}500{,}625} - \dfrac{121y^2}{3{,}339{,}375} = 1$ (b) ≈248 mi

55. (a) Ellipse (b) Hyperbola (c) No curve

59. 15.9

61. $\dfrac{b^2c}{a} + ab \ln\left(\dfrac{a}{b+c}\right)$ where $c^2 = a^2 + b^2$

63. $(0, 4/\pi)$

EXERCISES 10.6 ■ PAGE 708

1. $r = \dfrac{4}{2 + \cos\theta}$ **3.** $r = \dfrac{6}{2 + 3\sin\theta}$

5. $r = \dfrac{8}{1 - \sin\theta}$ **7.** $r = \dfrac{4}{2 + \cos\theta}$

9. (a) $\frac{4}{5}$ (b) Ellipse (c) $y = -1$
(d)



11. (a) 1 (b) Parabola (c) $y = \frac{2}{3}$
(d)



13. (a) $\frac{1}{3}$ (b) Ellipse (c) $x = \frac{9}{2}$
(d)



Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

15. (a) 2 (b) Hyperbola (c) $x = -\frac{3}{8}$
(d)



17. (a) $2, y = -\frac{1}{2}$



(b) $r = \dfrac{1}{1 - 2\sin(\theta - 3\pi/4)}$



19. The ellipse is nearly circular when e is close to 0 and becomes more elongated as $e \to 1^-$. At $e = 1$, the curve becomes a parabola.



25. $r = \dfrac{2.26 \times 10^8}{1 + 0.093\cos\theta}$

27. 35.64 AU **29.** 7.0×10^7 km **31.** 3.6×10^8 km

CHAPTER 10 REVIEW ■ PAGE 709

True-False Quiz

1. False **3.** False **5.** True **7.** False **9.** True

Exercises

1. $x = y^2 - 8y + 12$ **3.** $y = 1/x$





5. $x = t, y = \sqrt{t}$; $x = t^4, y = t^2$; $x = \tan^2 t, y = \tan t, 0 \leq t < \pi/2$

7. (a)



(b) $\left(3\sqrt{2}, 3\pi/4\right)$, $\left(-3\sqrt{2}, 7\pi/4\right)$

$\left(-2, 2\sqrt{3}\right)$

9.



11.



13.



15.



17. $r = \dfrac{2}{\cos\theta + \sin\theta}$

19.



21. 2 **23.** −1

25. $\dfrac{1 + \sin t}{1 + \cos t}, \dfrac{1 + \cos t + \sin t}{(1 + \cos t)^3}$ **27.** $\left(\frac{11}{8}, \frac{3}{4}\right)$

29. Vertical tangent at $\left(\frac{3}{2}a, \pm\frac{1}{2}\sqrt{3}\,a\right), (-3a, 0)$; horizontal tangent at $(a, 0), \left(-\frac{1}{2}a, \pm\frac{3}{2}\sqrt{3}a\right)$



31. 18 **33.** $(2, \pm\pi/3)$ **35.** $\frac{1}{2}(\pi - 1)$

37. $2\left(5\sqrt{5} - 1\right)$

39. $\dfrac{2\sqrt{\pi^2 + 1} - \sqrt{4\pi^2 + 1}}{2\pi} + \ln\left(\dfrac{2\pi + \sqrt{4\pi^2 + 1}}{\pi + \sqrt{\pi^2 + 1}}\right)$

41. $471{,}295\pi/1024$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

43. All curves have the vertical asymptote $x = 1$. For $c < -1$, the curve bulges to the right. At $c = -1$, the curve is the line $x = 1$. For $-1 < c < 0$, it bulges to the left. At $c = 0$ there is a cusp at $(0, 0)$. For $c > 0$, there is a loop.

45. $(\pm 1, 0), (\pm 3, 0)$



47. $\left(-\frac{25}{24}, 3\right), (-1, 3)$



49. $\dfrac{x^2}{25} + \dfrac{y^2}{9} = 1$ **51.** $\dfrac{y^2}{72/5} - \dfrac{x^2}{8/5} = 1$

53. $\dfrac{x^2}{25} + \dfrac{(8y - 399)^2}{160{,}801} = 1$ **55.** $r = \dfrac{4}{3 + \cos\theta}$

57. (a) At $(0, 0)$ and $\left(\frac{3}{2}, \frac{3}{2}\right)$
(b) Horizontal tangents at $(0, 0)$ and $\left(\sqrt[3]{2}, \sqrt[3]{4}\right)$; vertical tangents at $(0, 0)$ and $\left(\sqrt[3]{4}, \sqrt[3]{2}\right)$
(d)



(g) $\frac{3}{2}$

PROBLEMS PLUS ■ PAGE 712

1. $\ln(\pi/2)$ **3.** $\left[-\frac{3}{4}\sqrt{3}, \frac{3}{4}\sqrt{3}\right] \times [-1, 2]$

CHAPTER 11

EXERCISES 11.1 ■ PAGE 724

Abbreviations: C, convergent; D, divergent

1. (a) A sequence is an ordered list of numbers. It can also be defined as a function whose domain is the set of positive integers.
(b) The terms a_n approach 8 as n becomes large.
(c) The terms a_n become large as n becomes large.

3. $1, \frac{4}{5}, \frac{3}{5}, \frac{8}{17}, \frac{5}{13}$ **5.** $\frac{1}{5}, -\frac{1}{25}, \frac{1}{125}, -\frac{1}{625}, \frac{1}{3125}$ **7.** $\frac{1}{2}, \frac{1}{6}, \frac{1}{24}, \frac{1}{120}, \frac{1}{720}$

9. 1, 2, 7, 32, 157 **11.** $2, \frac{2}{3}, \frac{2}{5}, \frac{2}{7}, \frac{2}{9}$ **13.** $a_n = 1/(2n - 1)$

15. $a_n = -3\left(-\frac{2}{3}\right)^{n-1}$ **17.** $a_n = (-1)^{n+1}\dfrac{n^2}{n + 1}$

19. 0.4286, 0.4615, 0.4737, 0.4800, 0.4839, 0.4865, 0.4884, 0.4898, 0.4909, 0.4918; yes; $\frac{1}{2}$

21. 0.5000, 1.2500, 0.8750, 1.0625, 0.9688, 1.0156, 0.9922, 1.0039, 0.9980, 1.0010; yes; 1

23. 1 **25.** 5 **27.** 1 **29.** 1 **31.** D **33.** 0
35. D **37.** 0 **39.** 0 **41.** 0 **43.** 0 **45.** 1
47. e^2 **49.** $\ln 2$ **51.** $\pi/2$ **53.** D **55.** D
57. 1 **59.** $\frac{1}{2}$ **61.** D **63.** 0
65. (a) 1060, 1123.60, 1191.02, 1262.48, 1338.23 (b) D
67. (a) $P_n = 1.08P_{n-1} - 300$ (b) 5734
69. $-1 < r < 1$
71. Convergent by the Monotonic Sequence Theorem; $5 \leqslant L < 8$
73. Decreasing; yes **75.** Not monotonic; no
77. Decreasing; yes
79. 2 **81.** $\frac{1}{2}\left(3 + \sqrt{5}\right)$ **83.** (b) $\frac{1}{2}\left(1 + \sqrt{5}\right)$
85. (a) 0 (b) 9, 11

EXERCISES 11.2 ■ PAGE 735

1. (a) A sequence is an ordered list of numbers whereas a series is the *sum* of a list of numbers.
(b) A series is convergent if the sequence of partial sums is a convergent sequence. A series is divergent if it is not convergent.
3. 2
5. 1, 1.125, 1.1620, 1.1777, 1.1857, 1.1903, 1.1932, 1.1952; C
7. 0.5, 1.3284, 2.4265, 3.7598, 5.3049, 7.0443, 8.9644, 11.0540; D

9. −2.40000, −1.92000, −2.01600, −1.99680, −2.00064, −1.99987, −2.00003, −1.99999, −2.00000, −2.00000; convergent, sum = −2



11. 0.44721, 1.15432, 1.98637, 2.88080, 3.80927, 4.75796, 5.71948, 6.68962, 7.66581, 8.64639; divergent



13. 0.29289, 0.42265, 0.50000, 0.55279, 0.59175, 0.62204, 0.64645, 0.66667, 0.68377, 0.69849; convergent, sum = 1



15. (a) C (b) D **17.** D **19.** $\frac{25}{3}$ **21.** 60 **23.** $\frac{1}{7}$
25. D **27.** D **29.** D **31.** $\frac{5}{2}$ **33.** D **35.** D
37. D **39.** D **41.** $e/(e - 1)$ **43.** $\frac{3}{2}$ **45.** $\frac{11}{6}$ **47.** $e - 1$
49. (b) 1 (c) 2 (d) All rational numbers with a terminating decimal representation, except 0.
51. $\frac{8}{9}$ **53.** $\frac{838}{333}$ **55.** 5063/3300

57. $-\dfrac{1}{5} < x < \dfrac{1}{5}; \dfrac{-5x}{1 + 5x}$ **59.** $-1 < x < 5; \dfrac{3}{5 - x}$

61. $x > 2$ or $x < -2; \dfrac{x}{x - 2}$ **63.** $x < 0; \dfrac{1}{1 - e^x}$

65. 1 **67.** $a_1 = 0, a_n = \dfrac{2}{n(n + 1)}$ for $n > 1$, sum = 1

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

69. (a) 157.875 mg; $\frac{3000}{19}(1 - 0.05^n)$ (b) 157.895 mg

71. (a) $S_n = \dfrac{D(1 - c^n)}{1 - c}$ (b) 5 **73.** $\frac{1}{2}\left(\sqrt{3} - 1\right)$

77. $\dfrac{1}{n(n + 1)}$ **79.** The series is divergent.

85. $\{s_n\}$ is bounded and increasing.

87. (a) $0, \frac{1}{9}, \frac{2}{9}, \frac{1}{3}, \frac{2}{3}, \frac{7}{9}, \frac{8}{9}, 1$

89. (a) $\frac{1}{2}, \frac{5}{6}, \frac{23}{24}, \frac{119}{120}$; $\dfrac{(n + 1)! - 1}{(n + 1)!}$ (c) 1

EXERCISES 11.3 ■ PAGE 744

1. C



3. D **5.** C **7.** D **9.** C **11.** C **13.** D
15. C **17.** C **19.** C **21.** D **23.** C **25.** C
27. f is neither positive nor decreasing.
29. $p > 1$ **31.** $p < -1$ **33.** $(1, \infty)$
35. (a) $\frac{9}{10}\pi^4$ (b) $\frac{1}{90}\pi^4 - \frac{17}{16}$
37. (a) 1.54977, error $\leqslant 0.1$ (b) 1.64522, error $\leqslant 0.005$
(c) 1.64522 compared to 1.64493 (d) $n > 1000$
39. 0.00145 **45.** $b < 1/e$

EXERCISES 11.4 ■ PAGE 750

1. (a) Nothing (b) C **3.** C **5.** D **7.** C **9.** D
11. C **13.** C **15.** D **17.** D **19.** D **21.** C
23. C **25.** D **27.** C **29.** C **31.** D
33. 1.249, error < 0.1 **35.** 0.0739, error $< 6.4 \times 10^{-8}$
45. Yes

EXERCISES 11.5 ■ PAGE 755

1. (a) A series whose terms are alternately positive and negative (b) $0 < b_{n+1} \leqslant b_n$ and $\lim_{n\to\infty} b_n = 0$, where $b_n = |a_n|$ (c) $|R_n| \leqslant b_{n+1}$
3. C **5.** C **7.** D **9.** C **11.** C **13.** D **15.** C
17. C **19.** D **21.** -0.5507 **23.** 5 **25.** 4
27. -0.4597 **29.** 0.0676 **31.** An underestimate
33. p is not a negative integer **35.** $\{b_n\}$ is not decreasing

EXERCISES 11.6 ■ PAGE 761

Abbreviations: AC, absolutely convergent; CC, conditionally convergent

1. (a) D (b) C (c) May converge or diverge
3. AC **5.** CC **7.** AC **9.** D **11.** AC **13.** AC
15. AC **17.** CC **19.** AC **21.** AC **23.** D **25.** AC
27. AC **29.** D **31.** D **33.** AC
35. (a) and (d)
39. (a) $\frac{661}{960} \approx 0.68854$, error < 0.00521
(b) $n \geqslant 11$, 0.693109
45. (b) $\displaystyle\sum_{n=2}^{\infty} \frac{(-1)^n}{n \ln n}$; $\displaystyle\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$

EXERCISES 11.7 ■ PAGE 764

1. C **3.** D **5.** C **7.** D **9.** C **11.** C
13. C **15.** C **17.** C **19.** C **21.** D **23.** D
25. C **27.** C **29.** C **31.** D
33. C **35.** D **37.** C

EXERCISES 11.8 ■ PAGE 769

1. A series of the form $\sum_{n=0}^{\infty} c_n(x - a)^n$, where x is a variable and a and the c_n's are constants
3. $1, (-1, 1)$ **5.** $1, [-1, 1)$
7. $\infty, (-\infty, \infty)$ **9.** $2, (-2, 2)$ **11.** $\frac{1}{3}, \left[-\frac{1}{3}, \frac{1}{3}\right]$
13. $4, (-4, 4]$ **15.** $1, [1, 3]$ **17.** $\frac{1}{3}, \left[-\frac{13}{3}, -\frac{11}{3}\right)$
19. $\infty, (-\infty, \infty)$ **21.** $b, (a - b, a + b)$ **23.** $0, \left\{\frac{1}{2}\right\}$
25. $\frac{1}{5}, \left[\frac{3}{5}, 1\right]$ **27.** $\infty, (-\infty, \infty)$ **29.** (a) Yes (b) No
31. k^k **33.** No
35. (a) $(-\infty, \infty)$
(b), (c)



37. $(-1, 1), f(x) = (1 + 2x)/(1 - x^2)$ **41.** 2

EXERCISES 11.9 ■ PAGE 775

1. 10 **3.** $\displaystyle\sum_{n=0}^{\infty} (-1)^n x^n, (-1, 1)$ **5.** $\displaystyle 2\sum_{n=0}^{\infty} \frac{1}{3^{n+1}} x^n, (-3, 3)$

7. $\displaystyle\sum_{n=0}^{\infty} (-1)^n \frac{1}{9^{n+1}} x^{2n+1}, (-3, 3)$ **9.** $\displaystyle 1 + 2\sum_{n=1}^{\infty} x^n, (-1, 1)$

11. $\displaystyle\sum_{n=0}^{\infty} \left[(-1)^{n+1} - \frac{1}{2^{n+1}}\right] x^n, (-1, 1)$

13. (a) $\displaystyle\sum_{n=0}^{\infty} (-1)^n (n + 1) x^n, R = 1$

(b) $\displaystyle\frac{1}{2}\sum_{n=0}^{\infty} (-1)^n (n + 2)(n + 1) x^n, R = 1$

(c) $\displaystyle\frac{1}{2}\sum_{n=2}^{\infty} (-1)^n n(n - 1) x^n, R = 1$

15. $\displaystyle \ln 5 - \sum_{n=1}^{\infty} \frac{x^n}{n5^n}, R = 5$

17. $\displaystyle\sum_{n=0}^{\infty} (-1)^n 4^n (n + 1) x^{n+1}, R = \frac{1}{4}$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

19. $\sum_{n=0}^{\infty}(2n+1)x^n$, $R=1$

21. $\sum_{n=0}^{\infty}(-1)^n\frac{1}{16^{n+1}}x^{2n+1}$, $R=4$



23. $\sum_{n=0}^{\infty}\frac{2x^{2n+1}}{2n+1}$, $R=1$



25. $C+\sum_{n=0}^{\infty}\frac{t^{8n+2}}{8n+2}$, $R=1$

27. $C+\sum_{n=1}^{\infty}(-1)^n\frac{x^{n+3}}{n(n+3)}$, $R=1$

29. 0.199989 **31.** 0.000983 **33.** 0.19740

35. (b) 0.920 **39.** $[-1,1]$, $[-1,1)$, $(-1,1)$

EXERCISES 11.10 ■ PAGE 789

1. $b_8=f^{(8)}(5)/8!$ **3.** $\sum_{n=0}^{\infty}(n+1)x^n$, $R=1$

5. $\sum_{n=0}^{\infty}(n+1)x^n$, $R=1$

7. $\sum_{n=0}^{\infty}(-1)^n\frac{\pi^{2n+1}}{(2n+1)!}x^{2n+1}$, $R=\infty$

9. $\sum_{n=0}^{\infty}\frac{(\ln 2)^n}{n!}x^n$, $R=\infty$ **11.** $\sum_{n=0}^{\infty}\frac{x^{2n+1}}{(2n+1)!}$, $R=\infty$

13. $-1-2(x-1)+3(x-1)^2+4(x-1)^3+(x-1)^4$, $R=\infty$

15. $\ln 2+\sum_{n=1}^{\infty}(-1)^{n+1}\frac{1}{n\,2^n}(x-2)^n$, $R=2$

17. $\sum_{n=0}^{\infty}\frac{2^n e^6}{n!}(x-3)^n$, $R=\infty$

19. $\sum_{n=0}^{\infty}(-1)^{n+1}\frac{1}{(2n)!}(x-\pi)^{2n}$, $R=\infty$

25. $1-\frac{1}{4}x-\sum_{n=2}^{\infty}\frac{3\cdot 7\cdot\cdots\cdot(4n-5)}{4^n\cdot n!}x^n$, $R=1$

27. $\sum_{n=0}^{\infty}(-1)^n\frac{(n+1)(n+2)}{2^{n+4}}x^n$, $R=2$

29. $\sum_{n=0}^{\infty}(-1)^n\frac{\pi^{2n+1}}{(2n+1)!}x^{2n+1}$, $R=\infty$

31. $\sum_{n=0}^{\infty}\frac{2^n+1}{n!}x^n$, $R=\infty$

33. $\sum_{n=0}^{\infty}(-1)^n\frac{1}{2^{2n}(2n)!}x^{4n+1}$, $R=\infty$

35. $\frac{1}{2}x+\sum_{n=1}^{\infty}(-1)^n\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{n!\,2^{3n+1}}x^{2n+1}$, $R=2$

37. $\sum_{n=1}^{\infty}(-1)^{n+1}\frac{2^{2n-1}}{(2n)!}x^{2n}$, $R=\infty$

39. $\sum_{n=0}^{\infty}(-1)^n\frac{1}{(2n)!}x^{4n}$, $R=\infty$



41. $\sum_{n=1}^{\infty}\frac{(-1)^{n-1}}{(n-1)!}x^n$, $R=\infty$



43. 0.99619

45. (a) $1+\sum_{n=1}^{\infty}\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{2^n n!}x^{2n}$

(b) $x+\sum_{n=1}^{\infty}\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{(2n+1)2^n n!}x^{2n+1}$

47. $C+\sum_{n=0}^{\infty}(-1)^n\frac{x^{6n+2}}{(6n+2)(2n)!}$, $R=\infty$

49. $C+\sum_{n=1}^{\infty}(-1)^n\frac{1}{2n\,(2n)!}x^{2n}$, $R=\infty$

51. 0.0059 **53.** 0.40102 **55.** $\frac{1}{2}$ **57.** $\frac{1}{120}$

59. $1-\frac{3}{2}x^2+\frac{25}{24}x^4$ **61.** $1+\frac{1}{6}x^2+\frac{7}{360}x^4$ **63.** e^{-x^4}

65. $\ln\frac{8}{5}$ **67.** $1/\sqrt{2}$ **69.** e^3-1

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

EXERCISES 11.11 ■ PAGE 798

1. (a) $T_0(x) = 1 = T_1(x)$, $T_2(x) = 1 - \frac{1}{2}x^2 = T_3(x)$, $T_4(x) = 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 = T_5(x)$, $T_6(x) = 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6$



(b)

x	f	$T_0 = T_1$	$T_2 = T_3$	$T_4 = T_5$	T_6
$\frac{\pi}{4}$	0.7071	1	0.6916	0.7074	0.7071
$\frac{\pi}{2}$	0	1	−0.2337	0.0200	−0.0009
π	−1	1	−3.9348	0.1239	−1.2114

(c) As n increases, $T_n(x)$ is a good approximation to $f(x)$ on a larger and larger interval.

3. $\frac{1}{2} - \frac{1}{4}(x-2) + \frac{1}{8}(x-2)^2 - \frac{1}{16}(x-2)^3$



5. $-\left(x - \frac{\pi}{2}\right) + \frac{1}{6}\left(x - \frac{\pi}{2}\right)^3$



7. $(x-1) - \frac{1}{2}(x-1)^2 + \frac{1}{3}(x-1)^3$



9. $x - 2x^2 + 2x^3$



11. $T_5(x) = 1 - 2\left(x - \frac{\pi}{4}\right) + 2\left(x - \frac{\pi}{4}\right)^2 - \frac{8}{3}\left(x - \frac{\pi}{4}\right)^3 + \frac{10}{3}\left(x - \frac{\pi}{4}\right)^4 - \frac{64}{15}\left(x - \frac{\pi}{4}\right)^5$



13. (a) $2 + \frac{1}{4}(x-4) - \frac{1}{64}(x-4)^2$ (b) 1.5625×10^{-5}

15. (a) $1 + \frac{2}{3}(x-1) - \frac{1}{9}(x-1)^2 + \frac{4}{81}(x-1)^3$ (b) 0.000097

17. (a) $1 + \frac{1}{2}x^2$ (b) 0.0014

19. (a) $1 + x^2$ (b) 0.00006 **21.** (a) $x^2 - \frac{1}{6}x^4$ (b) 0.042

23. 0.17365 **25.** Four **27.** $-1.037 < x < 1.037$

29. $-0.86 < x < 0.86$ **31.** 21 m, no

37. (c) They differ by about 8×10^{-9} km.

CHAPTER 11 REVIEW ■ PAGE 802

True-False Quiz

1. False **3.** True **5.** False **7.** False **9.** False
11. True **13.** True **15.** False **17.** True
19. True **21.** True

Exercises

1. $\frac{1}{2}$ **3.** D **5.** 0 **7.** e^{12} **9.** 2 **11.** C **13.** C
15. D **17.** C **19.** C **21.** C **23.** CC **25.** AC

27. $\frac{1}{11}$ **29.** $\pi/4$ **31.** e^{-e} **35.** 0.9721

37. 0.18976224, error $< 6.4 \times 10^{-7}$

41. 4, $[-6, 2)$ **43.** 0.5, [2.5, 3.5)

45. $\frac{1}{2}\sum_{n=0}^{\infty}(-1)^n\left[\frac{1}{(2n)!}\left(x - \frac{\pi}{6}\right)^{2n} + \frac{\sqrt{3}}{(2n+1)!}\left(x - \frac{\pi}{6}\right)^{2n+1}\right]$

47. $\sum_{n=0}^{\infty}(-1)^n x^{n+2}$, $R = 1$ **49.** $\ln 4 - \sum_{n=1}^{\infty}\frac{x^n}{n4^n}$, $R = 4$

51. $\sum_{n=0}^{\infty}(-1)^n\frac{x^{8n+4}}{(2n+1)!}$, $R = \infty$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

53. $\frac{1}{2} + \sum_{n=1}^{\infty} \frac{1 \cdot 5 \cdot 9 \cdot \cdots \cdot (4n-3)}{n!\,2^{6n+1}} x^n$, $R = 16$

55. $C + \ln|x| + \sum_{n=1}^{\infty} \frac{x^n}{n \cdot n!}$

57. (a) $1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3$

(b) 

(c) 0.000006

59. $-\frac{1}{6}$

PROBLEMS PLUS ■ PAGE 805

1. $15!/5! = 10{,}897{,}286{,}400$

3. (b) 0 if $x = 0$, $(1/x) - \cot x$ if $x \neq k\pi$, k an integer

5. (a) $s_n = 3 \cdot 4^n$, $l_n = 1/3^n$, $p_n = 4^n/3^{n-1}$ (c) $\frac{2}{5}\sqrt{3}$

9. $(-1, 1)$, $\dfrac{x^3 + 4x^2 + x}{(1-x)^4}$

11. $\ln \frac{1}{2}$ **13.** (a) $\frac{250}{101}\pi(e^{-(n-1)\pi/5} - e^{-n\pi/5})$ (b) $\frac{250}{101}\pi$

19. $\dfrac{\pi}{2\sqrt{3}} - 1$

21. $-\left(\dfrac{\pi}{2} - \pi k\right)^2$ where k is a positive integer

CHAPTER 12

EXERCISES 12.1 ■ PAGE 814

1. $(4, 0, -3)$ **3.** C; A

5. A vertical plane that intersects the xy-plane in the line $y = 2 - x$, $z = 0$



7. (a) $|PQ| = 6$, $|QR| = 2\sqrt{10}$, $|RP| = 6$; isosceles triangle

9. (a) No (b) Yes

11. $(x+3)^2 + (y-2)^2 + (z-5)^2 = 16$;
$(y-2)^2 + (z-5)^2 = 7$, $x = 0$ (a circle)

13. $(x-3)^2 + (y-8)^2 + (z-1)^2 = 30$

15. $(1, 2, -4)$, 6 **17.** $(2, 0, -6)$, $9/\sqrt{2}$

19. (b) $\frac{5}{2}, \frac{1}{2}\sqrt{94}, \frac{1}{2}\sqrt{85}$

21. (a) $(x-2)^2 + (y+3)^2 + (z-6)^2 = 36$
(b) $(x-2)^2 + (y+3)^2 + (z-6)^2 = 4$
(c) $(x-2)^2 + (y+3)^2 + (z-6)^2 = 9$

23. A plane parallel to the yz-plane and 5 units in front of it

25. A half-space consisting of all points to the left of the plane $y = 8$

27. All points on or between the horizontal planes $z = 0$ and $z = 6$

29. All points on a circle with radius 2 with center on the z-axis that is contained in the plane $z = -1$

31. All points on or inside a sphere with radius $\sqrt{3}$ and center O

33. All points on or inside a circular cylinder of radius 3 with axis the y-axis

35. $0 < x < 5$ **37.** $r^2 < x^2 + y^2 + z^2 < R^2$

39. (a) $(2, 1, 4)$ (b) 

41. $14x - 6y - 10z = 9$, a plane perpendicular to AB

43. $2\sqrt{3} - 3$

EXERCISES 12.2 ■ PAGE 822

1. (a) Scalar (b) Vector (c) Vector (d) Scalar

3. $\overrightarrow{AB} = \overrightarrow{DC}$, $\overrightarrow{DA} = \overrightarrow{CB}$, $\overrightarrow{DE} = \overrightarrow{EB}$, $\overrightarrow{EA} = \overrightarrow{CE}$

5. (a) 

(b) 

(c)

(d) 

(e) 

(f) 

7. $\mathbf{c} = \frac{1}{2}\mathbf{a} + \frac{1}{2}\mathbf{b}$, $\mathbf{d} = \frac{1}{2}\mathbf{b} - \frac{1}{2}\mathbf{a}$

9. $\mathbf{a} = \langle 4, 1 \rangle$



11. $\mathbf{a} = \langle 3, -1 \rangle$



13. $\mathbf{a} = \langle 2, 0, -2 \rangle$



15. $\langle 5, 2 \rangle$



Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

17. $\langle 3, 8, 1 \rangle$



19. $\langle 2, -18 \rangle$, $\langle 1, -42 \rangle$, 13, 10
21. $-\mathbf{i} + \mathbf{j} + 2\mathbf{k}$, $-4\mathbf{i} + \mathbf{j} + 9\mathbf{k}$, $\sqrt{14}$, $\sqrt{82}$
23. $-\dfrac{3}{\sqrt{58}}\mathbf{i} + \dfrac{7}{\sqrt{58}}\mathbf{j}$ **25.** $\frac{8}{9}\mathbf{i} - \frac{1}{9}\mathbf{j} + \frac{4}{9}\mathbf{k}$ **27.** 60°
29. $\langle 2, 2\sqrt{3} \rangle$ **31.** ≈45.96 ft/s, ≈38.57 ft/s
33. $100\sqrt{7} \approx 264.6$ N, ≈139.1°
35. $\sqrt{493} \approx 22.2$ mi/h, N8°W
37. $\mathbf{T}_1 = -196\,\mathbf{i} + 3.92\,\mathbf{j}$, $\mathbf{T}_2 = 196\,\mathbf{i} + 3.92\,\mathbf{j}$
39. (a) At an angle of 43.4° from the bank, toward upstream
(b) 20.2 min
41. $\pm(\mathbf{i} + 4\,\mathbf{j})/\sqrt{17}$ **43.** 0
45. (a), (b)



(d) $s = \frac{9}{7}$, $t = \frac{11}{7}$
47. A sphere with radius 1, centered at (x_0, y_0, z_0)

EXERCISES 12.3 ■ PAGE 830

1. (b), (c), (d) are meaningful **3.** 14 **5.** 19 **7.** 1
9. −15 **11.** $\mathbf{u} \cdot \mathbf{v} = \frac{1}{2}$, $\mathbf{u} \cdot \mathbf{w} = -\frac{1}{2}$
15. $\cos^{-1}\left(\dfrac{1}{\sqrt{5}}\right) \approx 63°$ **17.** $\cos^{-1}\left(\dfrac{5}{\sqrt{1015}}\right) \approx 81°$
19. $\cos^{-1}\left(\dfrac{7}{\sqrt{130}}\right) \approx 52°$ **21.** 48°, 75°, 57°
23. (a) Neither (b) Orthogonal
(c) Orthogonal (d) Parallel
25. Yes **27.** $(\mathbf{i} - \mathbf{j} - \mathbf{k})/\sqrt{3}$ $\left[\text{or } (-\mathbf{i} + \mathbf{j} + \mathbf{k})/\sqrt{3}\right]$
29. 45° **31.** 0° at (0, 0), 8.1° at (1, 1)
33. $\frac{2}{3}, \frac{1}{3}, \frac{2}{3}$; 48°, 71°, 48°
35. $1/\sqrt{14}, -2/\sqrt{14}, -3/\sqrt{14}$; 74°, 122°, 143°
37. $1/\sqrt{3}, 1/\sqrt{3}, 1/\sqrt{3}$; 55°, 55°, 55° **39.** 4, $\left\langle -\frac{20}{13}, \frac{48}{13} \right\rangle$
41. $\frac{9}{7}$, $\left\langle \frac{27}{49}, \frac{54}{49}, -\frac{18}{49} \right\rangle$ **43.** $1/\sqrt{21}, \frac{2}{21}\,\mathbf{i} - \frac{1}{21}\,\mathbf{j} + \frac{4}{21}\,\mathbf{k}$
47. $\left\langle 0, 0, -2\sqrt{10} \right\rangle$ or any vector of the form
$\left\langle s, t, 3s - 2\sqrt{10} \right\rangle$, $s, t \in \mathbb{R}$
49. 144 J **51.** 2400 cos(40°) ≈ 1839 ft-lb
53. $\frac{13}{5}$ **55.** $\cos^{-1}\left(1/\sqrt{3}\right) \approx 55°$

EXERCISES 12.4 ■ PAGE 838

1. $16\,\mathbf{i} + 48\,\mathbf{k}$ **3.** $15\,\mathbf{i} - 3\,\mathbf{j} + 3\,\mathbf{k}$ **5.** $\frac{1}{2}\,\mathbf{i} - \mathbf{j} + \frac{3}{2}\,\mathbf{k}$
7. $(1 - t)\,\mathbf{i} + (t^3 - t^2)\,\mathbf{k}$ **9.** 0 **11.** $\mathbf{i} + \mathbf{j} + \mathbf{k}$
13. (a) Scalar (b) Meaningless (c) Vector
(d) Meaningless (e) Meaningless (f) Scalar
15. $96\sqrt{3}$; into the page **17.** $\langle -7, 10, 8 \rangle$, $\langle 7, -10, -8 \rangle$
19. $\left\langle -\dfrac{1}{3\sqrt{3}}, -\dfrac{1}{3\sqrt{3}}, \dfrac{5}{3\sqrt{3}} \right\rangle, \left\langle \dfrac{1}{3\sqrt{3}}, \dfrac{1}{3\sqrt{3}}, -\dfrac{5}{3\sqrt{3}} \right\rangle$
27. 16 **29.** (a) $\langle 0, 18, -9 \rangle$ (b) $\frac{9}{2}\sqrt{5}$
31. (a) $\langle 13, -14, 5 \rangle$ (b) $\frac{1}{2}\sqrt{390}$
33. 9 **35.** 16 **39.** $10.8 \sin 80° \approx 10.6$ N · m
41. ≈417 N **43.** 60°
45. (b) $\sqrt{97/3}$ **53.** (a) No (b) No (c) Yes

EXERCISES 12.5 ■ PAGE 848

1. (a) True (b) False (c) True (d) False (e) False
(f) True (g) False (h) True (i) True (j) False
(k) True
3. $\mathbf{r} = (2\,\mathbf{i} + 2.4\,\mathbf{j} + 3.5\,\mathbf{k}) + t(3\,\mathbf{i} + 2\,\mathbf{j} - \mathbf{k})$;
$x = 2 + 3t$, $y = 2.4 + 2t$, $z = 3.5 - t$
5. $\mathbf{r} = (\mathbf{i} + 6\mathbf{k}) + t(\mathbf{i} + 3\mathbf{j} + \mathbf{k})$;
$x = 1 + t$, $y = 3t$, $z = 6 + t$
7. $x = 2 + 2t$, $y = 1 + \frac{1}{2}t$, $z = -3 - 4t$;
$(x - 2)/2 = 2y - 2 = (z + 3)/(-4)$
9. $x = -8 + 11t$, $y = 1 - 3t$, $z = 4$; $\dfrac{x + 8}{11} = \dfrac{y - 1}{-3}$, $z = 4$
11. $x = 1 + t$, $y = -1 + 2t$, $z = 1 + t$;
$x - 1 = (y + 1)/2 = z - 1$
13. Yes
15. (a) $(x - 1)/(-1) = (y + 5)/2 = (z - 6)/(-3)$
(b) $(-1, -1, 0)$, $\left(-\frac{3}{2}, 0, -\frac{3}{2}\right)$, $(0, -3, 3)$
17. $\mathbf{r}(t) = (2\mathbf{i} - \mathbf{j} + 4\mathbf{k}) + t(2\mathbf{i} + 7\mathbf{j} - 3\mathbf{k})$, $0 \leqslant t \leqslant 1$
19. Skew **21.** $(4, -1, -5)$ **23.** $x - 2y + 5z = 0$
25. $x + 4y + z = 4$ **27.** $5x - y - z = 7$
29. $6x + 6y + 6z = 11$ **31.** $x + y + z = 2$
33. $-13x + 17y + 7z = -42$ **35.** $33x + 10y + 4z = 190$
37. $x - 2y + 4z = -1$ **39.** $3x - 8y - z = -38$
41.



43.



45. (2, 3, 5) **47.** (2, 3, 1) **49.** 1, 0, −1
51. Perpendicular **53.** Neither, $\cos^{-1}\left(\frac{1}{3}\right) \approx 70.5°$
55. Parallel
57. (a) $x = 1$, $y = -t$, $z = t$ (b) $\cos^{-1}\left(\dfrac{5}{3\sqrt{3}}\right) \approx 15.8°$
59. $x = 1$, $y - 2 = -z$ **61.** $x + 2y + z = 5$
63. $(x/a) + (y/b) + (z/c) = 1$
65. $x = 3t$, $y = 1 - t$, $z = 2 - 2t$
67. P_2 and P_3 are parallel, P_1 and P_4 are identical

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s).
Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

69. $\sqrt{61/14}$ **71.** $\frac{18}{7}$ **73.** $5/(2\sqrt{14})$ **77.** $1/\sqrt{6}$
79. $13/\sqrt{69}$

EXERCISES 12.6 ■ PAGE 856

1. (a) Parabola
(b) Parabolic cylinder with rulings parallel to the z-axis
(c) Parabolic cylinder with rulings parallel to the x-axis

3. Circular cylinder



5. Parabolic cylinder



7. Hyperbolic cylinder



9. (a) $x = k$, $y^2 - z^2 = 1 - k^2$, hyperbola ($k \neq \pm 1$);
$y = k$, $x^2 - z^2 = 1 - k^2$, hyperbola ($k \neq \pm 1$);
$z = k$, $x^2 + y^2 = 1 + k^2$, circle
(b) The hyperboloid is rotated so that it has axis the y-axis
(c) The hyperboloid is shifted one unit in the negative y-direction

11. Elliptic paraboloid with axis the x-axis



13. Elliptic cone with axis the x-axis



15. Hyperboloid of two sheets



17. Ellipsoid



19. Hyperbolic paraboloid



21. VII **23.** II **25.** VI **27.** VIII

29. $y^2 = x^2 + \dfrac{z^2}{9}$

Elliptic cone with axis the y-axis



31. $y = z^2 - \dfrac{x^2}{2}$

Hyperbolic paraboloid



33. $x^2 + \dfrac{(y-2)^2}{4} + (z-3)^2 = 1$

Ellipsoid with center
(0, 2, 3)



Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

35. $(y+1)^2 = (x-2)^2 + (z-1)^2$
Circular cone with vertex $(2, -1, 1)$ and axis parallel to the y-axis



37.



39.



41.



43. $y = x^2 + z^2$ **45.** $-4x = y^2 + z^2$, paraboloid

47. (a) $\dfrac{x^2}{(6378.137)^2} + \dfrac{y^2}{(6378.137)^2} + \dfrac{z^2}{(6356.523)^2} = 1$
(b) Circle (c) Ellipse

51.



CHAPTER 12 REVIEW ■ PAGE 858

True-False Quiz

1. False **3.** False **5.** True **7.** True **9.** True
11. True **13.** True **15.** False **17.** False
19. False **21.** True

Exercises

1. (a) $(x+1)^2 + (y-2)^2 + (z-1)^2 = 69$
(b) $(y-2)^2 + (z-1)^2 = 68,\ x = 0$
(c) Center $(4, -1, -3)$, radius 5
3. $\mathbf{u} \cdot \mathbf{v} = 3\sqrt{2};\ |\mathbf{u} \times \mathbf{v}| = 3\sqrt{2}$; out of the page
5. $-2, -4$ **7.** (a) 2 (b) -2 (c) -2 (d) 0
9. $\cos^{-1}\left(\frac{1}{3}\right) \approx 71°$ **11.** (a) $\langle 4, -3, 4 \rangle$ (b) $\sqrt{41}/2$
13. 166 N, 114 N
15. $x = 4 - 3t,\ y = -1 + 2t,\ z = 2 + 3t$
17. $x = -2 + 2t,\ y = 2 - t,\ z = 4 + 5t$
19. $-4x + 3y + z = -14$ **21.** (1, 4, 4) **23.** Skew
25. $x + y + z = 4$ **27.** $22/\sqrt{26}$

29. Plane



31. Cone



33. Hyperboloid of two sheets



35. Ellipsoid



37. $4x^2 + y^2 + z^2 = 16$

PROBLEMS PLUS ■ PAGE 861

1. $\left(\sqrt{3} - \frac{3}{2}\right)$ m
3. (a) $(x+1)/(-2c) = (y-c)/(c^2-1) = (z-c)/(c^2+1)$
(b) $x^2 + y^2 = t^2 + 1,\ z = t$ (c) $4\pi/3$
5. 20

CHAPTER 13

EXERCISES 13.1 ■ PAGE 869

1. $(-1, 2]$ **3.** $\mathbf{i} + \mathbf{j} + \mathbf{k}$ **5.** $\langle -1, \pi/2, 0 \rangle$

7.



9.



11.



13.



Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

15.







17. $\mathbf{r}(t) = \langle 2 + 4t, 2t, -2t \rangle$, $0 \leqslant t \leqslant 1$;
$x = 2 + 4t$, $y = 2t$, $z = -2t$, $0 \leqslant t \leqslant 1$

19. $\mathbf{r}(t) = \left\langle \frac{1}{2}t, -1 + \frac{4}{3}t, 1 - \frac{3}{4}t \right\rangle$, $0 \leqslant t \leqslant 1$;
$x = \frac{1}{2}t$, $y = -1 + \frac{4}{3}t$, $z = 1 - \frac{3}{4}t$, $0 \leqslant t \leqslant 1$

21. II **23.** V **25.** IV

27.



29. (0, 0, 0), (1, 0, 1)

31.



33.



35.



37.



41. $\mathbf{r}(t) = t\,\mathbf{i} + \frac{1}{2}(t^2 - 1)\,\mathbf{j} + \frac{1}{2}(t^2 + 1)\,\mathbf{k}$

43. $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j} + \cos 2t\,\mathbf{k}$, $0 \leqslant t \leqslant 2\pi$

45. $x = 2\cos t$, $y = 2\sin t$, $z = 4\cos^2 t$ **47.** Yes

EXERCISES 13.2 ■ PAGE 876

1. (a)



(b), (d)



(c) $\mathbf{r}'(4) = \lim_{h \to 0} \dfrac{\mathbf{r}(4 + h) - \mathbf{r}(4)}{h}$; $\mathbf{T}(4) = \dfrac{\mathbf{r}'(4)}{|\mathbf{r}'(4)|}$

3. (a), (c)



(b) $\mathbf{r}'(t) = \langle 1, 2t \rangle$

5. (a), (c)



(b) $\mathbf{r}'(t) = \cos t\,\mathbf{i} - 2\sin t\,\mathbf{j}$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

7. (a), (c)



(b) $\mathbf{r}'(t) = 2e^{2t}\,\mathbf{i} + e^{t}\,\mathbf{j}$

9. $\mathbf{r}'(t) = \langle t\cos t + \sin t,\ 2t,\ \cos 2t - 2t\sin 2t\rangle$

11. $\mathbf{r}'(t) = \mathbf{i} + \left(1/\sqrt{t}\right)\mathbf{k}$

13. $\mathbf{r}'(t) = 2te^{t^2}\,\mathbf{i} + [3/(1+3t)]\,\mathbf{k}$ **15.** $\mathbf{r}'(t) = \mathbf{b} + 2t\mathbf{c}$

17. $\left\langle \frac{1}{3}, \frac{2}{3}, \frac{2}{3}\right\rangle$ **19.** $\frac{3}{5}\,\mathbf{j} + \frac{4}{5}\,\mathbf{k}$

21. $\langle 1, 2t, 3t^2\rangle, \left\langle 1/\sqrt{14}, 2/\sqrt{14}, 3/\sqrt{14}\right\rangle, \langle 0, 2, 6t\rangle, \langle 6t^2, -6t, 2\rangle$

23. $x = 3 + t,\ y = 2t,\ z = 2 + 4t$

25. $x = 1 - t,\ y = t,\ z = 1 - t$

27. $\mathbf{r}(t) = (3 - 4t)\,\mathbf{i} + (4 + 3t)\,\mathbf{j} + (2 - 6t)\,\mathbf{k}$

29. $x = t,\ y = 1 - t,\ z = 2t$

31. $x = -\pi - t,\ y = \pi + t,\ z = -\pi t$

33. 66° **35.** $2\,\mathbf{i} - 4\,\mathbf{j} + 32\,\mathbf{k}$ **37.** $\mathbf{i} + \mathbf{j} + \mathbf{k}$

39. $\tan t\,\mathbf{i} + \frac{1}{8}(t^2+1)^4\mathbf{j} + \left(\frac{1}{3}t^3\ln t - \frac{1}{9}t^3\right)\mathbf{k} + \mathbf{C}$

41. $t^2\,\mathbf{i} + t^3\,\mathbf{j} + \left(\frac{2}{3}t^{3/2} - \frac{2}{3}\right)\mathbf{k}$

47. $2t\cos t + 2\sin t - 2\cos t\sin t$ **49.** 35

EXERCISES 13.3 ■ PAGE 884

1. $10\sqrt{10}$ **3.** $e - e^{-1}$ **5.** $\frac{1}{27}(13^{3/2} - 8)$ **7.** 18.6833

9. 1.2780 **11.** 42

13. $\mathbf{r}(t(s)) = \dfrac{2}{\sqrt{29}}s\,\mathbf{i} + \left(1 - \dfrac{3}{\sqrt{29}}s\right)\mathbf{j} + \left(5 + \dfrac{4}{\sqrt{29}}s\right)\mathbf{k}$

15. $(3\sin 1,\ 4,\ 3\cos 1)$

17. (a) $\left\langle 1/\sqrt{10}, \left(-3/\sqrt{10}\right)\sin t, \left(3/\sqrt{10}\right)\cos t\right\rangle$, $\langle 0, -\cos t, -\sin t\rangle$ (b) $\frac{3}{10}$

19. (a) $\dfrac{1}{e^{2t}+1}\left\langle \sqrt{2}e^{t}, e^{2t}, -1\right\rangle, \dfrac{1}{e^{2t}+1}\left\langle 1 - e^{2t}, \sqrt{2}e^{t}, \sqrt{2}e^{t}\right\rangle$

(b) $\sqrt{2}e^{2t}/(e^{2t}+1)^2$

21. $6t^2/(9t^4 + 4t^2)^{3/2}$ **23.** $\frac{4}{25}$ **25.** $\frac{1}{7}\sqrt{\frac{19}{14}}$

27. $12x^2/(1 + 16x^6)^{3/2}$ **29.** $e^x|x+2|/[1 + (xe^x + e^x)^2]^{3/2}$

31. $\left(-\frac{1}{2}\ln 2,\ 1/\sqrt{2}\right)$; approaches 0 **33.** (a) P (b) 1.3, 0.7

35.



37.





39. a is $y = f(x)$, b is $y = \kappa(x)$

41. $\kappa(t) = \dfrac{6\sqrt{4\cos^2 t - 12\cos t + 13}}{(17 - 12\cos t)^{3/2}}$



integer multiples of 2π

43. $6t^2/(4t^2 + 9t^4)^{3/2}$

45. $1/\left(\sqrt{2}e^{t}\right)$ **47.** $\left\langle \frac{2}{3}, \frac{2}{3}, \frac{1}{3}\right\rangle, \left\langle -\frac{1}{3}, \frac{2}{3}, -\frac{2}{3}\right\rangle, \left\langle -\frac{2}{3}, \frac{1}{3}, \frac{2}{3}\right\rangle$

49. $y = 6x + \pi,\ x + 6y = 6\pi$

51. $\left(x + \frac{5}{2}\right)^2 + y^2 = \frac{81}{4},\ x^2 + \left(y - \frac{5}{3}\right)^2 = \frac{16}{9}$



53. $(-1, -3, 1)$

55. $2x + y + 4z = 7,\ 6x - 8y - z = -3$

63. $2/(t^4 + 4t^2 + 1)$ **65.** 2.07×10^{10} Å ≈ 2 m

EXERCISES 13.4 ■ PAGE 894

1. (a) $1.8\,\mathbf{i} - 3.8\,\mathbf{j} - 0.7\,\mathbf{k}$, $2.0\,\mathbf{i} - 2.4\,\mathbf{j} - 0.6\,\mathbf{k}$, $2.8\,\mathbf{i} + 1.8\,\mathbf{j} - 0.3\,\mathbf{k}$, $2.8\,\mathbf{i} + 0.8\,\mathbf{j} - 0.4\,\mathbf{k}$

(b) $2.4\,\mathbf{i} - 0.8\,\mathbf{j} - 0.5\,\mathbf{k}$, 2.58

3. $\mathbf{v}(t) = \langle -t, 1\rangle$
$\mathbf{a}(t) = \langle -1, 0\rangle$
$|\mathbf{v}(t)| = \sqrt{t^2 + 1}$



5. $\mathbf{v}(t) = -3\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j}$
$\mathbf{a}(t) = -3\cos t\,\mathbf{i} - 2\sin t\,\mathbf{j}$
$|\mathbf{v}(t)| = \sqrt{5\sin^2 t + 4}$



7. $\mathbf{v}(t) = \mathbf{i} + 2t\,\mathbf{j}$
$\mathbf{a}(t) = 2\,\mathbf{j}$
$|\mathbf{v}(t)| = \sqrt{1 + 4t^2}$



Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

9. $\langle 2t+1, 2t-1, 3t^2\rangle$, $\langle 2, 2, 6t\rangle$, $\sqrt{9t^4+8t^2+2}$

11. $\sqrt{2}\,\mathbf{i}+e^t\,\mathbf{j}-e^{-t}\,\mathbf{k}$, $e^t\,\mathbf{j}+e^{-t}\,\mathbf{k}$, e^t+e^{-t}

13. $e^t[(\cos t-\sin t)\mathbf{i}+(\sin t+\cos t)\mathbf{j}+(t+1)\mathbf{k}]$, $e^t[-2\sin t\,\mathbf{i}+2\cos t\,\mathbf{j}+(t+2)\mathbf{k}]$, $e^t\sqrt{t^2+2t+3}$

15. $\mathbf{v}(t)=t\,\mathbf{i}+2t\,\mathbf{j}+\mathbf{k}$, $\mathbf{r}(t)=\left(\frac{1}{2}t^2+1\right)\mathbf{i}+t^2\,\mathbf{j}+t\,\mathbf{k}$

17. (a) $\mathbf{r}(t)=\left(\frac{1}{3}t^3+t\right)\mathbf{i}+(t-\sin t+1)\,\mathbf{j}+\left(\frac{1}{4}-\frac{1}{4}\cos 2t\right)\mathbf{k}$

(b)



19. $t=4$ **21.** $\mathbf{r}(t)=t\,\mathbf{i}-t\,\mathbf{j}+\frac{5}{2}t^2\,\mathbf{k}$, $|\mathbf{v}(t)|=\sqrt{25t^2+2}$

23. (a) ≈ 3535 m (b) ≈ 1531 m (c) 200 m/s

25. 30 m/s **27.** $\approx 10.2°$, $\approx 79.8°$

29. $13.0° < \theta < 36.0°$, $55.4° < \theta < 85.5°$

31. $(250, -50, 0)$; $10\sqrt{93}\approx 96.4$ ft/s

33. (a) 16 m (b) $\approx 23.6°$ upstream





35. The path is contained in a circle that lies in a plane perpendicular to $\mathbf{c}$ with center on a line through the origin in the direction of $\mathbf{c}$.

37. $6t$, 6 **39.** 0, 1 **41.** e^t-e^{-t}, $\sqrt{2}$

43. 4.5 cm/s^2, 9.0 cm/s^2 **45.** $t=1$

CHAPTER 13 REVIEW ■ PAGE 897

True-False Quiz

1. True **3.** False **5.** False **7.** False

9. True **11.** False **13.** True

Exercises

1. (a)



(b) $\mathbf{r}'(t)=\mathbf{i}-\pi\sin\pi t\,\mathbf{j}+\pi\cos\pi t\,\mathbf{k}$, $\mathbf{r}''(t)=-\pi^2\cos\pi t\,\mathbf{j}-\pi^2\sin\pi t\,\mathbf{k}$

3. $\mathbf{r}(t)=4\cos t\,\mathbf{i}+4\sin t\,\mathbf{j}+(5-4\cos t)\mathbf{k}$, $0\leqslant t\leqslant 2\pi$

5. $\frac{1}{3}\mathbf{i}-(2/\pi^2)\mathbf{j}+(2/\pi)\mathbf{k}$ **7.** 86.631 **9.** $\pi/2$

11. (a) $\langle t^2, t, 1\rangle/\sqrt{t^4+t^2+1}$

(b) $\langle t^3+2t, 1-t^4, -2t^3-t\rangle/\sqrt{t^8+5t^6+6t^4+5t^2+1}$

(c) $\sqrt{t^8+5t^6+6t^4+5t^2+1}/(t^4+t^2+1)^2$

13. $12/17^{3/2}$ **15.** $x-2y+2\pi=0$

17. $\mathbf{v}(t)=(1+\ln t)\,\mathbf{i}+\mathbf{j}-e^{-t}\,\mathbf{k}$, $|\mathbf{v}(t)|=\sqrt{2+2\ln t+(\ln t)^2+e^{-2t}}$, $\mathbf{a}(t)=(1/t)\mathbf{i}+e^{-t}\,\mathbf{k}$

19. (a) About 3.8 ft above the ground, 60.8 ft from the athlete

(b) ≈ 21.4 ft (c) ≈ 64.2 ft from the athlete

21. (c) $-2e^{-t}\,\mathbf{v}_d+e^{-t}\,\mathbf{R}$

23. (a) $\mathbf{v}=\omega R(-\sin\omega t\,\mathbf{i}+\cos\omega t\,\mathbf{j})$ (c) $\mathbf{a}=-\omega^2\mathbf{r}$

PROBLEMS PLUS ■ PAGE 900

1. (a) 90°, $v_0^2/(2g)$

3. (a) ≈ 0.94 ft to the right of the table's edge, ≈ 15 ft/s

(b) $\approx 7.6°$ (c) ≈ 2.13 ft to the right of the table's edge

5. 56°

7. $\mathbf{r}(u,v)=\mathbf{c}+u\,\mathbf{a}+v\,\mathbf{b}$ where $\mathbf{a}=\langle a_1, a_2, a_3\rangle$, $\mathbf{b}=\langle b_1, b_2, b_3\rangle$, $\mathbf{c}=\langle c_1, c_2, c_3\rangle$

CHAPTER 14

EXERCISES 14.1 ■ PAGE 912

1. (a) -27; a temperature of -15°C with wind blowing at 40 km/h feels equivalent to about -27°C without wind.

(b) When the temperature is -20°C, what wind speed gives a wind chill of -30°C? 20 km/h

(c) With a wind speed of 20 km/h, what temperature gives a wind chill of -49°C? -35°C

(d) A function of wind speed that gives wind-chill values when the temperature is -5°C

(e) A function of temperature that gives wind-chill values when the wind speed is 50 km/h

3. ≈ 94.2; the manufacturer's yearly production is valued at \$94.2 million when 120,000 labor hours are spent and \$20 million in capital is invested.

5. (a) ≈ 20.5; the surface area of a person 70 inches tall who weighs 160 pounds is approximately 20.5 square feet.

7. (a) 25; a 40-knot wind blowing in the open sea for 15 h will create waves about 25 ft high.

(b) $f(30, t)$ is a function of t giving the wave heights produced by 30-knot winds blowing for t hours.

(c) $f(v, 30)$ is a function of v giving the wave heights produced by winds of speed v blowing for 30 hours.

9. (a) 1 (b) $\mathbb{R}^2$ (c) $[-1, 1]$

11. (a) 3 (b) $\{(x,y,z)\mid x^2+y^2+z^2<4, x\geqslant 0, y\geqslant 0, z\geqslant 0\}$, interior of a sphere of radius 2, center the origin, in the first octant

13. $\{(x,y)\mid y\leqslant 2x\}$



15. $\left\{(x,y)\mid \frac{1}{9}x^2+y^2<1\right\}$, $(-\infty, \ln 9]$



Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

17. $\{(x, y) \mid -1 \leq x \leq 1, -1 \leq y \leq 1\}$



19. $\{(x, y) \mid y \geq x^2, x \neq \pm 1\}$



21. $\{(x, y, z) \mid x^2 + y^2 + z^2 \leq 1\}$



23. $z = 1 + y$, plane parallel to x-axis



25. $4x + 5y + z = 10$, plane



27. $z = y^2 + 1$, parabolic cylinder



29. $z = 9 - x^2 - 9y^2$, elliptic paraboloid



31. $z = \sqrt{4 - 4x^2 - y^2}$, top half of ellipsoid



33. ≈56, ≈35 **35.** 11°C, 19.5°C **37.** Steep; nearly flat

39.



41.



43. $(y - 2x)^2 = k$



45. $y = -\sqrt{x} + k$



47. $y = ke^{-x}$



49. $y^2 - x^2 = k^2$



Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

51. $x^2 + 9y^2 = k$



53.



55.



57.



59. (a) C (b) II **61.** (a) F (b) I

63. (a) B (b) VI **65.** Family of parallel planes

67. Family of circular cylinders with axis the x-axis ($k > 0$)

69. (a) Shift the graph of f upward 2 units

(b) Stretch the graph of f vertically by a factor of 2

(c) Reflect the graph of f about the xy-plane

(d) Reflect the graph of f about the xy-plane and then shift it upward 2 units

71.



f appears to have a maximum value of about 15. There are two local maximum points but no local minimum point.

73.



The function values approach 0 as x, y become large; as (x, y) approaches the origin, f approaches $\pm\infty$ or 0, depending on the direction of approach.

75. If $c = 0$, the graph is a cylindrical surface. For $c > 0$, the level curves are ellipses. The graph curves upward as we leave the origin, and the steepness increases as c increases. For $c < 0$, the level curves are hyperbolas. The graph curves upward in the y-direction and downward, approaching the xy-plane, in the x-direction giving a saddle-shaped appearance near (0, 0, 1).

77. $c = -2, 0, 2$ **79.** (b) $y = 0.75x + 0.01$

EXERCISES 14.2 ■ PAGE 923

1. Nothing; if f is continuous, $f(3, 1) = 6$ **3.** $-\frac{5}{2}$

5. 1 **7.** $\frac{2}{7}$ **9.** Does not exist **11.** Does not exist

13. 0 **15.** Does not exist **17.** 2

19. $\sqrt{3}$ **21.** Does not exist

23. The graph shows that the function approaches different numbers along different lines.

25. $h(x, y) = (2x + 3y - 6)^2 + \sqrt{2x + 3y - 6}$; $\{(x, y) \mid 2x + 3y \geqslant 6\}$

27. Along the line $y = x$ **29.** $\mathbb{R}^2$ **31.** $\{(x, y) \mid x^2 + y^2 \neq 1\}$

33. $\{(x, y) \mid x^2 + y^2 > 4\}$ **35.** $\{(x, y, z) \mid x^2 + y^2 + z^2 \leqslant 1\}$

37. $\{(x, y) \mid (x, y) \neq (0, 0)\}$ **39.** 0 **41.** -1

43.



f is continuous on $\mathbb{R}^2$

EXERCISES 14.3 ■ PAGE 935

1. (a) The rate of change of temperature as longitude varies, with latitude and time fixed; the rate of change as only latitude varies; the rate of change as only time varies.

(b) Positive, negative, positive

3. (a) $f_T(-15, 30) \approx 1.3$; for a temperature of -15°C and wind speed of 30 km/h, the wind-chill index rises by 1.3°C for each degree the temperature increases. $f_v(-15, 30) \approx -0.15$; for a temperature of -15°C and wind speed of 30 km/h, the wind-chill index decreases by 0.15°C for each km/h the wind speed increases.

(b) Positive, negative (c) 0

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

5. (a) Positive (b) Negative
7. (a) Positive (b) Negative
9. $c = f$, $b = f_x$, $a = f_y$
11. $f_x(1, 2) = -8$ = slope of C_1, $f_y(1, 2) = -4$ = slope of C_2



13.



$f(x, y) = x^2y^3$



$f_x(x, y) = 2xy^3$



$f_y(x, y) = 3x^2y^2$

15. $f_x(x, y) = -3y$, $f_y(x, y) = 5y^4 - 3x$
17. $f_x(x, t) = -\pi e^{-t} \sin \pi x$, $f_t(x, t) = -e^{-t} \cos \pi x$
19. $\partial z/\partial x = 20(2x + 3y)^9$, $\partial z/\partial y = 30(2x + 3y)^9$
21. $f_x(x, y) = 1/y$, $f_y(x, y) = -x/y^2$
23. $f_x(x, y) = \dfrac{(ad - bc)y}{(cx + dy)^2}$, $f_y(x, y) = \dfrac{(bc - ad)x}{(cx + dy)^2}$
25. $g_u(u, v) = 10uv(u^2v - v^3)^4$, $g_v(u, v) = 5(u^2 - 3v^2)(u^2v - v^3)^4$
27. $R_p(p, q) = \dfrac{q^2}{1 + p^2q^4}$, $R_q(p, q) = \dfrac{2pq}{1 + p^2q^4}$
29. $F_x(x, y) = \cos(e^x)$, $F_y(x, y) = -\cos(e^y)$
31. $f_x = z - 10xy^3z^4$, $f_y = -15x^2y^2z^4$, $f_z = x - 20x^2y^3z^3$
33. $\partial w/\partial x = 1/(x + 2y + 3z)$, $\partial w/\partial y = 2/(x + 2y + 3z)$, $\partial w/\partial z = 3/(x + 2y + 3z)$
35. $\partial u/\partial x = y \sin^{-1}(yz)$, $\partial u/\partial y = x \sin^{-1}(yz) + xyz/\sqrt{1 - y^2z^2}$, $\partial u/\partial z = xy^2/\sqrt{1 - y^2z^2}$
37. $h_x = 2xy \cos(z/t)$, $h_y = x^2 \cos(z/t)$, $h_z = (-x^2y/t) \sin(z/t)$, $h_t = (x^2yz/t^2) \sin(z/t)$
39. $\partial u/\partial x_i = x_i/\sqrt{x_1^2 + x_2^2 + \cdots + x_n^2}$
41. $\frac{1}{5}$ **43.** $\frac{1}{4}$ **45.** $f_x(x, y) = y^2 - 3x^2y$, $f_y(x, y) = 2xy - x^3$
47. $\dfrac{\partial z}{\partial x} = -\dfrac{x}{3z}$, $\dfrac{\partial z}{\partial y} = -\dfrac{2y}{3z}$
49. $\dfrac{\partial z}{\partial x} = \dfrac{yz}{e^z - xy}$, $\dfrac{\partial z}{\partial y} = \dfrac{xz}{e^z - xy}$
51. (a) $f'(x)$, $g'(y)$ (b) $f'(x + y)$, $f'(x + y)$
53. $f_{xx} = 6xy^5 + 24x^2y$, $f_{xy} = 15x^2y^4 + 8x^3 = f_{yx}$, $f_{yy} = 20x^3y^3$
55. $w_{uu} = v^2/(u^2 + v^2)^{3/2}$, $w_{uv} = -uv/(u^2 + v^2)^{3/2} = w_{vu}$, $w_{vv} = u^2/(u^2 + v^2)^{3/2}$
57. $z_{xx} = -2x/(1 + x^2)^2$, $z_{xy} = 0 = z_{yx}$, $z_{yy} = -2y/(1 + y^2)^2$
63. $24xy^2 - 6y$, $24x^2y - 6x$ **65.** $(2x^2y^2z^5 + 6xyz^3 + 2z)e^{xyz^2}$
67. $\theta e^{r\theta}(2 \sin \theta + \theta \cos \theta + r\theta \sin \theta)$ **69.** $4/(y + 2z)^3$, 0
71. $6yz^2$ **73.** ≈12.2, ≈16.8, ≈23.25 **83.** R^2/R_1^2
87. $\dfrac{\partial T}{\partial P} = \dfrac{V - nb}{nR}$, $\dfrac{\partial P}{\partial V} = \dfrac{2n^2a}{V^3} - \dfrac{nRT}{(V - nb)^2}$
93. No **95.** $x = 1 + t$, $y = 2$, $z = 2 - 2t$ **99.** -2
101. (a)



(b) $f_x(x, y) = \dfrac{x^4y + 4x^2y^3 - y^5}{(x^2 + y^2)^2}$, $f_y(x, y) = \dfrac{x^5 - 4x^3y^2 - xy^4}{(x^2 + y^2)^2}$
(c) 0, 0 (e) No, since f_{xy} and f_{yx} are not continuous.

EXERCISES 14.4 ■ PAGE 946

1. $z = -7x - 6y + 5$ **3.** $x + y - 2z = 0$
5. $x + y + z = 0$
7.



9.



11. $6x + 4y - 23$ **13.** $\frac{1}{9}x - \frac{2}{9}y + \frac{2}{3}$ **15.** $1 - \pi y$
19. 6.3 **21.** $\frac{3}{7}x + \frac{2}{7}y + \frac{6}{7}z$; 6.9914
23. $4T + H - 329$; 129°F
25. $dz = -2e^{-2x} \cos 2\pi t\, dx - 2\pi e^{-2x} \sin 2\pi t\, dt$
27. $dm = 5p^4q^3\, dp + 3p^5q^2\, dq$
29. $dR = \beta^2 \cos \gamma\, d\alpha + 2\alpha\beta \cos \gamma\, d\beta - \alpha\beta^2 \sin \gamma\, d\gamma$
31. $\Delta z = 0.9225$, $dz = 0.9$ **33.** 5.4 cm² **35.** 16 cm³
37. $\approx -0.0165mg$; decrease
39. $\frac{1}{17} \approx 0.059\ \Omega$ **41.** 2.3% **43.** $\varepsilon_1 = \Delta x$, $\varepsilon_2 = \Delta y$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

EXERCISES 14.5 ■ PAGE 954

1. $(2x + y)\cos t + (2y + x)e^t$

3. $[(x/t) - y\sin t]/\sqrt{1 + x^2 + y^2}$

5. $e^{y/z}[2t - (x/z) - (2xy/z^2)]$

7. $\partial z/\partial s = 2xy^3\cos t + 3x^2y^2\sin t$,
$\partial z/\partial t = -2sxy^3\sin t + 3sx^2y^2\cos t$

9. $\partial z/\partial s = t^2\cos\theta\cos\phi - 2st\sin\theta\sin\phi$,
$\partial z/\partial t = 2st\cos\theta\cos\phi - s^2\sin\theta\sin\phi$

11. $\dfrac{\partial z}{\partial s} = e^r\left(t\cos\theta - \dfrac{s}{\sqrt{s^2+t^2}}\sin\theta\right)$,
$\dfrac{\partial z}{\partial t} = e^r\left(s\cos\theta - \dfrac{t}{\sqrt{s^2+t^2}}\sin\theta\right)$

13. 62 **15.** 7, 2

17. $\dfrac{\partial u}{\partial r} = \dfrac{\partial u}{\partial x}\dfrac{\partial x}{\partial r} + \dfrac{\partial u}{\partial y}\dfrac{\partial y}{\partial r}$, $\dfrac{\partial u}{\partial s} = \dfrac{\partial u}{\partial x}\dfrac{\partial x}{\partial s} + \dfrac{\partial u}{\partial y}\dfrac{\partial y}{\partial s}$,
$\dfrac{\partial u}{\partial t} = \dfrac{\partial u}{\partial x}\dfrac{\partial x}{\partial t} + \dfrac{\partial u}{\partial y}\dfrac{\partial y}{\partial t}$

19. $\dfrac{\partial w}{\partial x} = \dfrac{\partial w}{\partial r}\dfrac{\partial r}{\partial x} + \dfrac{\partial w}{\partial s}\dfrac{\partial s}{\partial x} + \dfrac{\partial w}{\partial t}\dfrac{\partial t}{\partial x}$,
$\dfrac{\partial w}{\partial y} = \dfrac{\partial w}{\partial r}\dfrac{\partial r}{\partial y} + \dfrac{\partial w}{\partial s}\dfrac{\partial s}{\partial y} + \dfrac{\partial w}{\partial t}\dfrac{\partial t}{\partial y}$

21. 1582, 3164, −700 **23.** 2π, -2π

25. $\frac{5}{144}$, $-\frac{5}{96}$, $\frac{5}{144}$ **27.** $\dfrac{2x + y\sin x}{\cos x - 2y}$

29. $\dfrac{1 + x^4y^2 + y^2 + x^4y^4 - 2xy}{x^2 - 2xy - 2x^5y^3}$

31. $-\dfrac{x}{3z}, -\dfrac{2y}{3z}$ **33.** $\dfrac{yz}{e^z - xy}, \dfrac{xz}{e^z - xy}$

35. 2°C/s **37.** ≈ −0.33 m/s per minute

39. (a) 6 m³/s (b) 10 m²/s (c) 0 m/s

41. ≈ −0.27 L/s **43.** $-1/(12\sqrt{3})$ rad/s

45. (a) $\partial z/\partial r = (\partial z/\partial x)\cos\theta + (\partial z/\partial y)\sin\theta$,
$\partial z/\partial\theta = -(\partial z/\partial x)r\sin\theta + (\partial z/\partial y)r\cos\theta$

51. $4rs\,\partial^2z/\partial x^2 + (4r^2 + 4s^2)\partial^2z/\partial x\,\partial y + 4rs\,\partial^2z/\partial y^2 + 2\,\partial z/\partial y$

EXERCISES 14.6 ■ PAGE 967

1. ≈ −0.08 mb/km **3.** ≈ 0.778 **5.** $2 + \sqrt{3}/2$

7. (a) $\nabla f(x, y) = \langle 2\cos(2x + 3y), 3\cos(2x + 3y)\rangle$
(b) $\langle 2, 3\rangle$ (c) $\sqrt{3} - \frac{3}{2}$

9. (a) $\langle 2xyz - yz^3, x^2z - xz^3, x^2y - 3xyz^2\rangle$
(b) $\langle -3, 2, 2\rangle$ (c) $\frac{2}{5}$

11. $\dfrac{4 - 3\sqrt{3}}{10}$ **13.** $-8/\sqrt{10}$ **15.** $4/\sqrt{30}$

17. $\frac{23}{42}$ **19.** 2/5 **21.** $\sqrt{65}$, $\langle 1, 8\rangle$

23. 1, $\langle 0, 1\rangle$ **25.** 1, $\langle 3, 6, -2\rangle$

27. (b) $\langle -12, 92\rangle$

29. All points on the line $y = x + 1$

31. (a) $-40/(3\sqrt{3})$

33. (a) $32/\sqrt{3}$ (b) $\langle 38, 6, 12\rangle$ (c) $2\sqrt{406}$

35. $\frac{327}{13}$ **39.** $\frac{774}{25}$

41. (a) $x + y + z = 11$ (b) $x - 3 = y - 3 = z - 5$

43. (a) $2x + 3y + 12z = 24$ (b) $\dfrac{x-3}{2} = \dfrac{y-2}{3} = \dfrac{z-1}{12}$

45. (a) $x + y + z = 1$ (b) $x = y = z - 1$

47.



49. $\langle 2, 3\rangle$, $2x + 3y = 12$



55. No **59.** $\left(-\frac{5}{4}, -\frac{5}{4}, \frac{25}{8}\right)$

63. $x = -1 - 10t$, $y = 1 - 16t$, $z = 2 - 12t$

67. If $\mathbf{u} = \langle a, b\rangle$ and $\mathbf{v} = \langle c, d\rangle$, then $af_x + bf_y$ and $cf_x + df_y$ are known, so we solve linear equations for f_x and f_y.

EXERCISES 14.7 ■ PAGE 977

1. (a) f has a local minimum at (1, 1).
(b) f has a saddle point at (1, 1).

3. Local minimum at (1, 1), saddle point at (0, 0)

5. Minimum $f\left(\frac{1}{3}, -\frac{2}{3}\right) = -\frac{1}{3}$

7. Saddle points at (1, 1), (−1, −1)

9. Maximum $f(0, 0) = 2$, minimum $f(0, 4) = -30$,
saddle points at (2, 2), (−2, 2)

11. Minimum $f(2, 1) = -8$, saddle point at (0, 0)

13. None **15.** Minimum $f(0, 0) = 0$, saddle points at $(\pm 1, 0)$

17. Minima $f(0, 1) = f(\pi, -1) = f(2\pi, 1) = -1$,
saddle points at $(\pi/2, 0)$, $(3\pi/2, 0)$

21. Minima $f(1, \pm 1) = 3$, $f(-1, \pm 1) = 3$

23. Maximum $f(\pi/3, \pi/3) = 3\sqrt{3}/2$,
minimum $f(5\pi/3, 5\pi/3) = -3\sqrt{3}/2$, saddle point at (π, π)

25. Minima $f(0, -0.794) \approx -1.191$, $f(\pm 1.592, 1.267) \approx -1.310$,
saddle points $(\pm 0.720, 0.259)$,
lowest points $(\pm 1.592, 1.267, -1.310)$

27. Maximum $f(0.170, -1.215) \approx 3.197$,
minima $f(-1.301, 0.549) \approx -3.145$, $f(1.131, 0.549) \approx -0.701$,
saddle points $(-1.301, -1.215)$, $(0.170, 0.549)$, $(1.131, -1.215)$,
no highest or lowest point

29. Maximum $f(0, \pm 2) = 4$, minimum $f(1, 0) = -1$

31. Maximum $f(\pm 1, 1) = 7$, minimum $f(0, 0) = 4$

33. Maximum $f(3, 0) = 83$, minimum $f(1, 1) = 0$

35. Maximum $f(1, 0) = 2$, minimum $f(-1, 0) = -2$

37.



Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

39. $2/\sqrt{3}$ **41.** $(2, 1, \sqrt{5}), (2, 1, -\sqrt{5})$ **43.** $\frac{100}{3}, \frac{100}{3}, \frac{100}{3}$
45. $8r^3/(3\sqrt{3})$ **47.** $\frac{4}{3}$ **49.** Cube, edge length $c/12$
51. Square base of side 40 cm, height 20 cm **53.** $L^3/(3\sqrt{3})$

EXERCISES 14.8 ■ PAGE 987

1. ≈59, 30
3. No maximum, minimum $f(1, 1) = f(-1, -1) = 2$
5. Maximum $f(0, \pm 1) = 1$, minimum $f(\pm 2, 0) = -4$
7. Maximum $f(2, 2, 1) = 9$, minimum $f(-2, -2, -1) = -9$
9. Maximum $2/\sqrt{3}$, minimum $-2/\sqrt{3}$
11. Maximum $\sqrt{3}$, minimum 1
13. Maximum $f(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}) = 2$,
minimum $f(-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}) = -2$
15. Maximum $f(1, \sqrt{2}, -\sqrt{2}) = 1 + 2\sqrt{2}$,
minimum $f(1, -\sqrt{2}, \sqrt{2}) = 1 - 2\sqrt{2}$
17. Maximum $\frac{3}{2}$, minimum $\frac{1}{2}$
19. Maximum $f(3/\sqrt{2}, -3/\sqrt{2}) = 9 + 12\sqrt{2}$,
minimum $f(-2, 2) = -8$
21. Maximum $f(\pm 1/\sqrt{2}, \mp 1/(2\sqrt{2})) = e^{1/4}$,
minimum $f(\pm 1/\sqrt{2}, \pm 1/(2\sqrt{2})) = e^{-1/4}$
29–41. See Exercises 39–53 in Section 14.7.
43. Nearest $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, farthest $(-1, -1, 2)$
45. Maximum ≈9.7938, minimum ≈−5.3506
47. (a) c/n (b) When $x_1 = x_2 = \cdots = x_n$

CHAPTER 14 REVIEW ■ PAGE 991

True-False Quiz

1. True **3.** False **5.** False **7.** True **9.** False
11. True

Exercises

1. $\{(x, y) \mid y > -x - 1\}$



3.



5.



7.



9. $\frac{2}{3}$
11. (a) ≈3.5°C/m, −3.0°C/m (b) ≈0.35°C/m by Equation 14.6.9 (Definition 14.6.2 gives ≈1.1°C/m.)
(c) −0.25
13. $f_x = 32xy(5y^3 + 2x^2y)^7$, $f_y = (16x^2 + 120y^2)(5y^3 + 2x^2y)^7$
15. $F_\alpha = \dfrac{2\alpha^3}{\alpha^2 + \beta^2} + 2\alpha \ln(\alpha^2 + \beta^2)$, $F_\beta = \dfrac{2\alpha^2\beta}{\alpha^2 + \beta^2}$
17. $S_u = \arctan(v\sqrt{w})$, $S_v = \dfrac{u\sqrt{w}}{1 + v^2w}$, $S_w = \dfrac{uv}{2\sqrt{w}(1 + v^2w)}$
19. $f_{xx} = 24x$, $f_{xy} = -2y = f_{yx}$, $f_{yy} = -2x$
21. $f_{xx} = k(k-1)x^{k-2}y^lz^m$, $f_{xy} = klx^{k-1}y^{l-1}z^m = f_{yx}$,
$f_{xz} = kmx^{k-1}y^lz^{m-1} = f_{zx}$, $f_{yy} = l(l-1)x^ky^{l-2}z^m$,
$f_{yz} = lmx^ky^{l-1}z^{m-1} = f_{zy}$, $f_{zz} = m(m-1)x^ky^lz^{m-2}$
25. (a) $z = 8x + 4y + 1$ (b) $\dfrac{x-1}{8} = \dfrac{y+2}{4} = \dfrac{z-1}{-1}$
27. (a) $2x - 2y - 3z = 3$ (b) $\dfrac{x-2}{4} = \dfrac{y+1}{-4} = \dfrac{z-1}{-6}$
29. (a) $x + 2y + 5z = 0$
(b) $x = 2 + t$, $y = -1 + 2t$, $z = 5t$
31. $(2, \frac{1}{2}, -1), (-2, -\frac{1}{2}, 1)$
33. $60x + \frac{24}{5}y + \frac{32}{5}z - 120$; 38.656
35. $2xy^3(1 + 6p) + 3x^2y^2(pe^p + e^p) + 4z^3(p\cos p + \sin p)$
37. −47, 108
43. $\langle 2xe^{yz^2}, x^2z^2e^{yz^2}, 2x^2yze^{yz^2}\rangle$ **45.** $-\frac{4}{5}$
47. $\sqrt{145}/2, \langle 4, \frac{9}{2}\rangle$ **49.** ≈$\frac{5}{8}$ knot/mi
51. Minimum $f(-4, 1) = -11$
53. Maximum $f(1, 1) = 1$; saddle points (0, 0), (0, 3), (3, 0)
55. Maximum $f(1, 2) = 4$, minimum $f(2, 4) = -64$
57. Maximum $f(-1, 0) = 2$, minima $f(1, \pm 1) = -3$,
saddle points $(-1, \pm 1)$, $(1, 0)$
59. Maximum $f(\pm\sqrt{2/3}, 1/\sqrt{3}) = 2/(3\sqrt{3})$,
minimum $f(\pm\sqrt{2/3}, -1/\sqrt{3}) = -2/(3\sqrt{3})$
61. Maximum 1, minimum −1
63. $(\pm 3^{-1/4}, 3^{-1/4}\sqrt{2}, \pm 3^{1/4}), (\pm 3^{-1/4}, -3^{-1/4}\sqrt{2}, \pm 3^{1/4})$
65. $P(2 - \sqrt{3})$, $P(3 - \sqrt{3})/6$, $P(2\sqrt{3} - 3)/3$

PROBLEMS PLUS ■ PAGE 995

1. L^2W^2, $\frac{1}{4}L^2W^2$ **3.** (a) $x = w/3$, base $= w/3$ (b) Yes
7. $\sqrt{3/2}, 3/\sqrt{2}$

CHAPTER 15

EXERCISES 15.1 ■ PAGE 1005

1. (a) 288 (b) 144 **3.** (a) 0.990 (b) 1.151
5. (a) 4 (b) −8 **7.** $U < V < L$
9. (a) ≈248 (b) ≈15.5 **11.** 60 **13.** 3
15. 1.141606, 1.143191, 1.143535, 1.143617, 1.143637, 1.143642

EXERCISES 15.2 ■ PAGE 1011

1. $500y^3$, $3x^2$ **3.** 222 **5.** $32(e^4 - 1)$ **7.** 18
9. $\frac{21}{2}\ln 2$ **11.** $\frac{31}{30}$ **13.** π **15.** 0

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

17. $9 \ln 2$ **19.** $\frac{1}{2}\left(\sqrt{3}-1\right)-\frac{1}{12}\pi$ **21.** $\frac{1}{2}e^{-6}+\frac{5}{2}$

23.



25. 51 **27.** $\frac{166}{27}$ **29.** 2 **31.** $\frac{64}{3}$

33. $21e-57$



35. $\frac{5}{6}$ **37.** 0

39. Fubini's Theorem does not apply. The integrand has an infinite discontinuity at the origin.

EXERCISES 15.3 ■ PAGE 1019

1. 32 **3.** $\frac{3}{10}$ **5.** $\frac{1}{3}\sin 1$ **7.** $\frac{4}{3}$ **9.** π

11. (a)



(b)



13. Type I: $D=\{(x,y)\mid 0\leqslant x\leqslant 1,\ 0\leqslant y\leqslant x\}$, type II: $D=\{(x,y)\mid 0\leqslant y\leqslant 1,\ y\leqslant x\leqslant 1\}$; $\frac{1}{3}$

15. $\int_0^1\int_{-\sqrt{x}}^{\sqrt{x}} y\,dy\,dx+\int_1^4\int_{x-2}^{\sqrt{x}} y\,dy\,dx=\int_{-1}^{2}\int_{y^2}^{y+2} y\,dx\,dy=\frac{9}{4}$

17. $\frac{1}{2}(1-\cos 1)$ **19.** $\frac{11}{3}$ **21.** 0 **23.** $\frac{17}{60}$ **25.** $\frac{31}{8}$

27. 6 **29.** $\frac{128}{15}$ **31.** $\frac{1}{3}$ **33.** 0, 1.213; 0.713 **35.** $\frac{64}{3}$

37.



39. 13,984,735,616/14,549,535

41. $\pi/2$

43. $\int_0^1\int_x^1 f(x,y)\,dy\,dx$



45. $\int_0^1\int_0^{\cos^{-1}y} f(x,y)\,dx\,dy$



47. $\int_0^{\ln 2}\int_{e^y}^{2} f(x,y)\,dx\,dy$



49. $\frac{1}{6}(e^9-1)$ **51.** $\frac{1}{3}\ln 9$ **53.** $\frac{1}{3}\left(2\sqrt{2}-1\right)$ **55.** 1

57. $(\pi/16)e^{-1/16}\leqslant\iint_Q e^{-(x^2+y^2)^2}\,dA\leqslant\pi/16$ **59.** $\frac{3}{4}$ **63.** 9π

65. $a^2b+\frac{3}{2}ab^2$ **67.** $\pi a^2 b$

EXERCISES 15.4 ■ PAGE 1026

1. $\int_0^{3\pi/2}\int_0^4 f(r\cos\theta, r\sin\theta)r\,dr\,d\theta$ **3.** $\int_{-1}^{1}\int_0^{(x+1)/2} f(x,y)\,dy\,dx$

5.



$3\pi/4$

7. $\frac{1250}{3}$ **9.** $(\pi/4)(\cos 1-\cos 9)$

11. $(\pi/2)(1-e^{-4})$ **13.** $\frac{3}{64}\pi^2$ **15.** $\pi/12$

17. $\frac{\pi}{3}+\frac{\sqrt{3}}{2}$ **19.** $\frac{16}{3}\pi$ **21.** $\frac{4}{3}\pi$ **23.** $\frac{4}{3}\pi a^3$

25. $(2\pi/3)\left[1-\left(1/\sqrt{2}\right)\right]$ **27.** $(8\pi/3)\left(64-24\sqrt{3}\right)$

29. $\frac{1}{2}\pi(1-\cos 9)$ **31.** $2\sqrt{2}/3$ **33.** 4.5951

35. $1800\pi\ \text{ft}^3$ **37.** $2/(a+b)$ **39.** $\frac{15}{16}$

41. (a) $\sqrt{\pi}/4$ (b) $\sqrt{\pi}/2$

EXERCISES 15.5 ■ PAGE 1036

1. 285 C **3.** $42k, \left(2, \frac{85}{28}\right)$ **5.** $6, \left(\frac{3}{4}, \frac{3}{2}\right)$ **7.** $\frac{8}{15}k, \left(0, \frac{4}{7}\right)$

9. $L/4, (L/2, 16/(9\pi))$ **11.** $\left(\frac{3}{8}, 3\pi/16\right)$ **13.** $(0, 45/(14\pi))$

15. $(2a/5, 2a/5)$ if vertex is (0, 0) and sides are along positive axes

17. $\frac{64}{315}k, \frac{8}{105}k, \frac{88}{315}k$

19. $7ka^6/180, 7ka^6/180, 7ka^6/90$ if vertex is (0, 0) and sides are along positive axes

21. $\rho bh^3/3, \rho b^3h/3; b/\sqrt{3}, h/\sqrt{3}$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

23. $\rho a^4\pi/16,\ \rho a^4\pi/16;\ a/2,\ a/2$

25. $m = 3\pi/64,\ (\bar{x}, \bar{y}) = \left(\dfrac{16384\sqrt{2}}{10395\pi}, 0\right),$
$I_x = \dfrac{5\pi}{384} - \dfrac{4}{105},\ I_y = \dfrac{5\pi}{384} + \dfrac{4}{105},\ I_0 = \dfrac{5\pi}{192}$

27. (a) $\frac{1}{2}$ (b) 0.375 (c) $\frac{5}{48} \approx 0.1042$

29. (b) (i) $e^{-0.2} \approx 0.8187$
(ii) $1 + e^{-1.8} - e^{-0.8} - e^{-1} \approx 0.3481$ (c) 2, 5

31. (a) ≈ 0.500 (b) ≈ 0.632

33. (a) $\iint_D k\left[1 - \frac{1}{20}\sqrt{(x - x_0)^2 + (y - y_0)^2}\right] dA$, where D is the disk with radius 10 mi centered at the center of the city
(b) $200\pi k/3 \approx 209k$, $200\left(\pi/2 - \frac{8}{9}\right)k \approx 136k$, on the edge

EXERCISES 15.6 ■ PAGE 1040

1. $15\sqrt{26}$ **3.** $3\sqrt{14}$ **5.** $12\sin^{-1}\left(\frac{2}{3}\right)$
7. $(\pi/6)\left(17\sqrt{17} - 5\sqrt{5}\right)$ **9.** $(2\pi/3)\left(2\sqrt{2} - 1\right)$
11. $a^2(\pi - 2)$ **13.** 13.9783 **15.** (a) ≈ 1.83 (b) ≈ 1.8616
17. $\frac{45}{8}\sqrt{14} + \frac{15}{16}\ln\left[\left(11\sqrt{5} + 3\sqrt{70}\right)/\left(3\sqrt{5} + \sqrt{70}\right)\right]$
19. 3.3213 **23.** $(\pi/6)\left(101\sqrt{101} - 1\right)$

EXERCISES 15.7 ■ PAGE 1049

1. $\frac{27}{4}$ **3.** $\frac{16}{15}$ **5.** $\frac{5}{3}$ **7.** $-\frac{1}{3}$ **9.** $\frac{27}{2}$ **11.** $9\pi/8$
13. $\frac{65}{28}$ **15.** $\frac{1}{60}$ **17.** $16\pi/3$ **19.** $\frac{16}{3}$ **21.** $\frac{8}{15}$

23. (a) $\int_0^1 \int_0^x \int_0^{\sqrt{1-y^2}} dz\, dy\, dx$ (b) $\frac{1}{4}\pi - \frac{1}{3}$

25. 0.985

27.



29. $\int_{-2}^{2} \int_0^{4-x^2} \int_{-\sqrt{4-x^2-y}/2}^{\sqrt{4-x^2-y}/2} f(x, y, z)\, dz\, dy\, dx$
$= \int_0^4 \int_{-\sqrt{4-y}}^{\sqrt{4-y}} \int_{-\sqrt{4-x^2-y}/2}^{\sqrt{4-x^2-y}/2} f(x, y, z)\, dz\, dx\, dy$
$= \int_{-1}^{1} \int_0^{4-4z^2} \int_{-\sqrt{4-y-4z^2}}^{\sqrt{4-y-4z^2}} f(x, y, z)\, dx\, dy\, dz$
$= \int_0^4 \int_{-\sqrt{4-y}/2}^{\sqrt{4-y}/2} \int_{-\sqrt{4-y-4z^2}}^{\sqrt{4-y-4z^2}} f(x, y, z)\, dx\, dz\, dy$
$= \int_{-2}^{2} \int_{-\sqrt{4-x^2}/2}^{\sqrt{4-x^2}/2} \int_0^{4-x^2-4z^2} f(x, y, z)\, dy\, dz\, dx$
$= \int_{-1}^{1} \int_{-\sqrt{4-4z^2}}^{\sqrt{4-4z^2}} \int_0^{4-x^2-4z^2} f(x, y, z)\, dy\, dx\, dz$

31. $\int_{-2}^{2} \int_{x^2}^{4} \int_0^{2-y/2} f(x, y, z)\, dz\, dy\, dx$
$= \int_0^4 \int_{-\sqrt{y}}^{\sqrt{y}} \int_0^{2-y/2} f(x, y, z)\, dz\, dx\, dy$
$= \int_0^2 \int_0^{4-2z} \int_{-\sqrt{y}}^{\sqrt{y}} f(x, y, z)\, dx\, dy\, dz$
$= \int_0^4 \int_0^{2-y/2} \int_{-\sqrt{y}}^{\sqrt{y}} f(x, y, z)\, dx\, dz\, dy$
$= \int_{-2}^{2} \int_0^{2-x^2/2} \int_{x^2}^{4-2z} f(x, y, z)\, dy\, dz\, dx$
$= \int_0^2 \int_{-\sqrt{4-2z}}^{\sqrt{4-2z}} \int_{x^2}^{4-2z} f(x, y, z)\, dy\, dx\, dz$

33. $\int_0^1 \int_{\sqrt{x}}^1 \int_0^{1-y} f(x, y, z)\, dz\, dy\, dx = \int_0^1 \int_0^{y^2} \int_0^{1-y} f(x, y, z)\, dz\, dx\, dy$
$= \int_0^1 \int_0^{1-z} \int_0^{y^2} f(x, y, z)\, dx\, dy\, dz = \int_0^1 \int_0^{1-y} \int_0^{y^2} f(x, y, z)\, dx\, dz\, dy$
$= \int_0^1 \int_0^{1-\sqrt{x}} \int_{\sqrt{x}}^{1-z} f(x, y, z)\, dy\, dz\, dx = \int_0^1 \int_0^{(1-z)^2} \int_{\sqrt{x}}^{1-z} f(x, y, z)\, dy\, dx\, dz$

35. $\int_0^1 \int_y^1 \int_0^y f(x, y, z)\, dz\, dx\, dy = \int_0^1 \int_0^x \int_0^y f(x, y, z)\, dz\, dy\, dx$
$= \int_0^1 \int_z^1 \int_y^1 f(x, y, z)\, dx\, dy\, dz = \int_0^1 \int_0^y \int_y^1 f(x, y, z)\, dx\, dz\, dy$
$= \int_0^1 \int_0^x \int_z^x f(x, y, z)\, dy\, dz\, dx = \int_0^1 \int_z^1 \int_z^x f(x, y, z)\, dy\, dx\, dz$

37. 64π **39.** $\frac{79}{30}, \left(\frac{358}{553}, \frac{33}{79}, \frac{571}{553}\right)$

41. $a^5,\ (7a/12,\ 7a/12,\ 7a/12)$

43. $I_x = I_y = I_z = \frac{2}{3}kL^5$ **45.** $\frac{1}{2}\pi khka^4$

47. (a) $m = \int_{-1}^{1} \int_{x^2}^{1} \int_0^{1-y} \sqrt{x^2 + y^2}\, dz\, dy\, dx$
(b) $(\bar{x}, \bar{y}, \bar{z})$, where
$\bar{x} = (1/m) \int_{-1}^{1} \int_{x^2}^{1} \int_0^{1-y} x\sqrt{x^2 + y^2}\, dz\, dy\, dx$
$\bar{y} = (1/m) \int_{-1}^{1} \int_{x^2}^{1} \int_0^{1-y} y\sqrt{x^2 + y^2}\, dz\, dy\, dx$
$\bar{z} = (1/m) \int_{-1}^{1} \int_{x^2}^{1} \int_0^{1-y} z\sqrt{x^2 + y^2}\, dz\, dy\, dx$
(c) $\int_{-1}^{1} \int_{x^2}^{1} \int_0^{1-y} (x^2 + y^2)^{3/2}\, dz\, dy\, dx$

49. (a) $\frac{3}{32}\pi + \frac{11}{24}$

(b) $\left(\dfrac{28}{9\pi + 44}, \dfrac{30\pi + 128}{45\pi + 220}, \dfrac{45\pi + 208}{135\pi + 660}\right)$

(c) $\frac{1}{240}(68 + 15\pi)$

51. (a) $\frac{1}{8}$ (b) $\frac{1}{64}$ (c) $\frac{1}{5760}$ **53.** $L^3/8$

55. (a) The region bounded by the ellipsoid $x^2 + 2y^2 + 3z^2 = 1$
(b) $4\sqrt{6}\pi/45$

EXERCISES 15.8 ■ PAGE 1055

1. (a)



$\left(2, 2\sqrt{3}, -2\right)$

(b)



$(0, -2, 1)$

3. (a) $\left(\sqrt{2}, 3\pi/4, 1\right)$ (b) $(4, 2\pi/3, 3)$

5. Vertical half-plane through the z-axis

7. Circular paraboloid

9. (a) $z^2 = 1 + r\cos\theta - r^2$ (b) $z = r^2\cos 2\theta$

11.



13. Cylindrical coordinates: $6 \leq r \leq 7,\ 0 \leq \theta \leq 2\pi,\ 0 \leq z \leq 20$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

15. 4π



17. 384π **19.** $\frac{8}{3}\pi + \frac{128}{15}$ **21.** $2\pi/5$ **23.** $\frac{4}{3}\pi\left(\sqrt{2} - 1\right)$
25. (a) 162π (b) (0, 0, 15)
27. $\pi Ka^2/8$, (0, 0, $2a/3$) **29.** 0
31. (a) $\iiint_C h(P)g(P)\,dV$, where C is the cone
(b) $\approx 3.1 \times 10^{19}$ ft-lb

EXERCISES 15.9 ■ PAGE 1061

1. (a)



$\left(\frac{3}{2}, \frac{3\sqrt{3}}{2}, 3\sqrt{3}\right)$

(b)



$\left(0, \frac{3\sqrt{2}}{2}, -\frac{3\sqrt{2}}{2}\right)$

3. (a) $(2, 3\pi/2, \pi/2)$ (b) $(2, 3\pi/4, 3\pi/4)$
5. Half-cone **7.** Sphere, radius $\frac{1}{2}$, center $\left(0, \frac{1}{2}, 0\right)$
9. (a) $\cos^2\phi = \sin^2\phi$ (b) $\rho^2(\sin^2\phi\cos^2\theta + \cos^2\phi) = 9$
11.



13.



15. $0 \leq \phi \leq \pi/4$, $0 \leq \rho \leq \cos\phi$

17. $(9\pi/4)\left(2 - \sqrt{3}\right)$



19. $\int_0^{\pi/2}\int_0^3\int_0^2 f(r\cos\theta, r\sin\theta, z)\,r\,dz\,dr\,d\theta$
21. $312{,}500\pi/7$ **23.** $1688\pi/15$ **25.** $\pi/8$
27. $\left(\sqrt{3} - 1\right)\pi a^3/3$ **29.** (a) 10π (b) (0, 0, 2.1)
31. (a) $\left(0, 0, \frac{7}{12}\right)$ (b) $11K\pi/960$
33. (a) $\left(0, 0, \frac{3}{8}a\right)$ (b) $4K\pi a^5/15$
35. $\frac{1}{3}\pi\left(2 - \sqrt{2}\right)$, $\left(0, 0, 3/\left[8\left(2 - \sqrt{2}\right)\right]\right)$
37. $5\pi/6$ **39.** $\left(4\sqrt{2} - 5\right)/15$ **41.** $4096\pi/21$
43.



45. $136\pi/99$

EXERCISES 15.10 ■ PAGE 1071

1. 16 **3.** $\sin^2\theta - \cos^2\theta$ **5.** 0
7. The parallelogram with vertices (0, 0), (6, 3), (12, 1), (6, −2)
9. The region bounded by the line $y = 1$, the y-axis, and $y = \sqrt{x}$
11. $x = \frac{1}{3}(v - u)$, $y = \frac{1}{3}(u + 2v)$ is one possible transformation, where $S = \{(u, v) \mid -1 \leq u \leq 1, 1 \leq v \leq 3\}$
13. $x = u\cos v$, $y = u\sin v$ is one possible transformation, where $S = \left\{(u, v) \mid 1 \leq u \leq \sqrt{2}, 0 \leq v \leq \pi/2\right\}$
15. -3 **17.** 6π **19.** $2\ln 3$
21. (a) $\frac{4}{3}\pi abc$ (b) 1.083×10^{12} km^3 (c) $\frac{4}{15}\pi(a^2 + b^2)abck$
23. $\frac{8}{5}\ln 8$ **25.** $\frac{3}{2}\sin 1$ **27.** $e - e^{-1}$

CHAPTER 15 REVIEW ■ PAGE 1073

True-False Quiz

1. True **3.** True **5.** True **7.** True **9.** False

Exercises

1. ≈ 64.0 **3.** $4e^2 - 4e + 3$ **5.** $\frac{1}{2}\sin 1$ **7.** $\frac{2}{3}$
9. $\int_0^{\pi}\int_2^4 f(r\cos\theta, r\sin\theta)\,r\,dr\,d\theta$
11. The region inside the loop of the four-leaved rose $r = \sin 2\theta$ in the first quadrant
13. $\frac{1}{2}\sin 1$ **15.** $\frac{1}{2}e^6 - \frac{7}{2}$ **17.** $\frac{1}{4}\ln 2$ **19.** 8
21. $81\pi/5$ **23.** $\frac{81}{2}$ **25.** $\pi/96$ **27.** $\frac{64}{15}$
29. 176 **31.** $\frac{2}{3}$ **33.** $2ma^3/9$
35. (a) $\frac{1}{4}$ (b) $\left(\frac{1}{3}, \frac{8}{15}\right)$
(c) $I_x = \frac{1}{12}$, $I_y = \frac{1}{24}$; $\bar{\bar{y}} = 1/\sqrt{3}$, $\bar{\bar{x}} = 1/\sqrt{6}$
37. (a) $(0, 0, h/4)$ (b) $\pi a^4 h/10$
39. $\ln\left(\sqrt{2} + \sqrt{3}\right) + \sqrt{2}/3$ **41.** $\frac{486}{5}$ **43.** 0.0512

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

45. (a) $\frac{1}{15}$ (b) $\frac{1}{3}$ (c) $\frac{1}{45}$

47. $\int_0^1 \int_0^{1-z} \int_{-\sqrt{y}}^{\sqrt{y}} f(x, y, z)\, dx\, dy\, dz$ **49.** $-\ln 2$ **51.** 0

PROBLEMS PLUS ■ PAGE 1077

1. 30 **3.** $\frac{1}{2}\sin 1$ **7.** (b) 0.90

13. $abc\pi\left(\frac{2}{3} - \frac{8}{9\sqrt{3}}\right)$

CHAPTER 16

EXERCISES 16.1 ■ PAGE 1085

1.



3.



5.



7.



9.



11. IV **13.** I **15.** IV **17.** III

19.



The line $y = 2x$

21. $\nabla f(x, y) = (xy + 1)e^{xy}\,\mathbf{i} + x^2 e^{xy}\,\mathbf{j}$

23. $\nabla f(x, y, z) = \dfrac{x}{\sqrt{x^2 + y^2 + z^2}}\,\mathbf{i} + \dfrac{y}{\sqrt{x^2 + y^2 + z^2}}\,\mathbf{j} + \dfrac{z}{\sqrt{x^2 + y^2 + z^2}}\,\mathbf{k}$

25. $\nabla f(x, y) = 2x\,\mathbf{i} - \mathbf{j}$



27.



29. III **31.** II **33.** (2.04, 1.03)

35. (a)



(b) $y = 1/x,\ x > 0$

$y = C/x$

EXERCISES 16.2 ■ PAGE 1096

1. $\frac{1}{54}(145^{3/2} - 1)$ **3.** 1638.4 **5.** $\frac{243}{8}$ **7.** $\frac{5}{2}$

9. $\sqrt{5}\pi$ **11.** $\frac{1}{12}\sqrt{14}(e^6 - 1)$ **13.** $\frac{2}{5}(e - 1)$ **15.** $\frac{35}{3}$

17. (a) Positive (b) Negative **19.** 45

21. $\frac{6}{5} - \cos 1 - \sin 1$ **23.** 1.9633 **25.** 15.0074

27. $3\pi + \frac{2}{3}$



Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

29. (a) $\frac{11}{8} - 1/e$ (b)



31. $\frac{172{,}704}{5{,}632{,}705}\sqrt{2}(1 - e^{-14\pi})$ **33.** $2\pi k$, $(4/\pi, 0)$

35. (a) $\bar{x} = (1/m)\int_C x\rho(x, y, z)\, ds$,
$\bar{y} = (1/m)\int_C y\rho(x, y, z)\, ds$,
$\bar{z} = (1/m)\int_C z\rho(x, y, z)\, ds$, where $m = \int_C \rho(x, y, z)\, ds$
(b) $(0, 0, 3\pi)$

37. $I_x = k\left(\frac{1}{2}\pi - \frac{4}{3}\right)$, $I_y = k\left(\frac{1}{2}\pi - \frac{2}{3}\right)$ **39.** $2\pi^2$ **41.** $\frac{7}{3}$

43. (a) $2ma\,\mathbf{i} + 6mbt\,\mathbf{j}$, $0 \leq t \leq 1$ (b) $2ma^2 + \frac{9}{2}mb^2$

45. $\approx 1.67 \times 10^4$ ft-lb **47.** (b) Yes **51.** ≈ 22 J

EXERCISES 16.3 ■ PAGE 1106

1. 40 **3.** $f(x, y) = x^2 - 3xy + 2y^2 - 8y + K$

5. Not conservative **7.** $f(x, y) = ye^x + x \sin y + K$

9. $f(x, y) = x \ln y + x^2y^3 + K$

11. (b) 16 **13.** (a) $f(x, y) = \frac{1}{2}x^2y^2$ (b) 2

15. (a) $f(x, y, z) = xyz + z^2$ (b) 77

17. (a) $f(x, y, z) = ye^{xz}$ (b) 4 **19.** $4/e$

21. It doesn't matter which curve is chosen.

23. 30 **25.** No **27.** Conservative

31. (a) Yes (b) Yes (c) Yes

33. (a) No (b) Yes (c) Yes

EXERCISES 16.4 ■ PAGE 1113

1. 8π **3.** $\frac{2}{3}$ **5.** 12 **7.** $\frac{1}{3}$ **9.** -24π **11.** $-\frac{16}{3}$

13. 4π **15.** $-8e + 48e^{-1}$ **17.** $-\frac{1}{12}$ **19.** 3π **21.** (c) $\frac{9}{2}$

23. $(4a/3\pi, 4a/3\pi)$ if the region is the portion of the disk $x^2 + y^2 = a^2$ in the first quadrant

27. 0

EXERCISES 16.5 ■ PAGE 1121

1. (a) $\mathbf{0}$ (b) 3

3. (a) $ze^x\,\mathbf{i} + (xye^z - yze^x)\,\mathbf{j} - xe^z\,\mathbf{k}$ (b) $y(e^z + e^x)$

5. (a) $\mathbf{0}$ (b) $2/\sqrt{x^2 + y^2 + z^2}$

7. (a) $\langle -e^y \cos z, -e^z \cos x, -e^x \cos y \rangle$
(b) $e^x \sin y + e^y \sin z + e^z \sin x$

9. (a) Negative (b) curl $\mathbf{F} = \mathbf{0}$

11. (a) Zero (b) curl $\mathbf{F}$ points in the negative z-direction

13. $f(x, y, z) = xy^2z^3 + K$ **15.** Not conservative

17. $f(x, y, z) = xe^{yz} + K$ **19.** No

EXERCISES 16.6 ■ PAGE 1132

1. P: no; Q: yes

3. Plane through (0, 3, 1) containing vectors $\langle 1, 0, 4 \rangle$, $\langle 1, -1, 5 \rangle$

5. Hyperbolic paraboloid

7.



9.



11.



13. IV **15.** II **17.** III

19. $x = u, y = v - u, z = -v$

21. $y = y, z = z, x = \sqrt{1 + y^2 + \frac{1}{4}z^2}$

23. $x = 2 \sin\phi \cos\theta$, $y = 2 \sin\phi \sin\theta$,
$z = 2 \cos\phi$, $0 \leq \phi \leq \pi/4$, $0 \leq \theta \leq 2\pi$
$\left[\text{or } x = x, y = y, z = \sqrt{4 - x^2 - y^2}, x^2 + y^2 \leq 2\right]$

25. $x = x, y = 4 \cos\theta, z = 4 \sin\theta$, $0 \leq x \leq 5$, $0 \leq \theta \leq 2\pi$

29. $x = x, y = e^{-x} \cos\theta$,
$z = e^{-x} \sin\theta$, $0 \leq x \leq 3$,
$0 \leq \theta \leq 2\pi$



31. (a) Direction reverses (b) Number of coils doubles

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

33. $3x - y + 3z = 3$ **35.** $\frac{\sqrt{3}}{2}x - \frac{1}{2}y + z = \frac{\pi}{3}$

37. $-x + 2z = 1$ **39.** $3\sqrt{14}$ **41.** $\sqrt{14}\,\pi$

43. $\frac{4}{15}(3^{5/2} - 2^{7/2} + 1)$ **45.** $(2\pi/3)(2\sqrt{2} - 1)$

47. $\frac{1}{2}\sqrt{21} + \frac{17}{4}\left[\ln\left(2 + \sqrt{21}\right) - \ln\sqrt{17}\right]$ **49.** 4

51. $A(S) \leqslant \sqrt{3}\,\pi R^2$ **53.** 13.9783

55. (a) 24.2055 (b) 24.2476

57. $\frac{45}{8}\sqrt{14} + \frac{15}{16}\ln\left[\left(11\sqrt{5} + 3\sqrt{70}\right)/\left(3\sqrt{5} + \sqrt{70}\right)\right]$

59. (b)



(c) $\int_0^{2\pi}\int_0^{\pi}\sqrt{36\sin^4u\cos^2v + 9\sin^4u\sin^2v + 4\cos^2u\sin^2u}\,du\,dv$

61. 4π **63.** $2a^2(\pi - 2)$

EXERCISES 16.7 ■ PAGE 1144

1. 49.09 **3.** 900π **5.** $11\sqrt{14}$ **7.** $\frac{2}{3}(2\sqrt{2} - 1)$

9. $171\sqrt{14}$ **11.** $\sqrt{21}/3$ **13.** $364\sqrt{2}\,\pi/3$

15. $(\pi/60)(391\sqrt{17} + 1)$ **17.** 16π **19.** 12 **21.** 4

23. $\frac{713}{180}$ **25.** $-\frac{4}{3}\pi$ **27.** 0 **29.** 48 **31.** $2\pi + \frac{8}{3}$

33. 4.5822 **35.** 3.4895

37. $\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_D [P(\partial h/\partial x) - Q + R(\partial h/\partial z)]\,dA$, where D = projection of S on xz-plane

39. $(0, 0, a/2)$

41. (a) $I_z = \iint_S (x^2 + y^2)\rho(x, y, z)\,dS$ (b) $4329\sqrt{2}\,\pi/5$

43. 0 kg/s **45.** $\frac{8}{3}\pi a^3\varepsilon_0$ **47.** 1248π

EXERCISES 16.8 ■ PAGE 1151

3. 0 **5.** 0 **7.** -1 **9.** 80π

11. (a) $81\pi/2$ (b)



(c) $x = 3\cos t$, $y = 3\sin t$, $z = 1 - 3(\cos t + \sin t)$, $0 \leqslant t \leqslant 2\pi$



17. 3

EXERCISES 16.9 ■ PAGE 1157

5. $\frac{9}{2}$ **7.** $9\pi/2$ **9.** 0 **11.** $32\pi/3$ **13.** 2π

15. $341\sqrt{2}/60 + \frac{81}{20}\arcsin\left(\sqrt{3}/3\right)$

17. $13\pi/20$ **19.** Negative at P_1, positive at P_2

21. div $\mathbf{F} > 0$ in quadrants I, II; div $\mathbf{F} < 0$ in quadrants III, IV

CHAPTER 16 REVIEW ■ PAGE 1160

True-False Quiz

1. False **3.** True **5.** False
7. False **9.** True **11.** True

Exercises

1. (a) Negative (b) Positive **3.** $6\sqrt{10}$ **5.** $\frac{4}{15}$

7. $\frac{110}{3}$ **9.** $\frac{11}{12} - 4/e$ **11.** $f(x, y) = e^y + xe^{xy}$ **13.** 0

17. -8π **25.** $\frac{1}{6}(27 - 5\sqrt{5})$ **27.** $(\pi/60)(391\sqrt{17} + 1)$

29. $-64\pi/3$ **33.** $-\frac{1}{2}$ **37.** -4 **39.** 21

CHAPTER 17

EXERCISES 17.1 ■ PAGE 1172

1. $y = c_1e^{3x} + c_2e^{-2x}$ **3.** $y = c_1\cos 4x + c_2\sin 4x$

5. $y = c_1e^{2x/3} + c_2xe^{2x/3}$ **7.** $y = c_1 + c_2e^{x/2}$

9. $y = e^{2x}(c_1\cos 3x + c_2\sin 3x)$

11. $y = c_1e^{(\sqrt{3}-1)t/2} + c_2e^{-(\sqrt{3}+1)t/2}$

13. $P = e^{-t}\left[c_1\cos\left(\frac{1}{10}t\right) + c_2\sin\left(\frac{1}{10}t\right)\right]$

15.



All solutions approach either 0 or $\pm\infty$ as $x \to \pm\infty$.

17. $y = 3e^{2x} - e^{4x}$ **19.** $y = e^{-2x/3} + \frac{2}{3}xe^{-2x/3}$

21. $y = e^{3x}(2\cos x - 3\sin x)$

23. $y = \frac{1}{7}e^{4x-4} - \frac{1}{7}e^{3-3x}$ **25.** $y = 5\cos 2x + 3\sin 2x$

27. $y = 2e^{-2x} - 2xe^{-2x}$ **29.** $y = \dfrac{e - 2}{e - 1} + \dfrac{e^x}{e - 1}$

31. No solution

33. (b) $\lambda = n^2\pi^2/L^2$, n a positive integer; $y = C\sin(n\pi x/L)$

35. (a) $b - a \neq n\pi$, n any integer

(b) $b - a = n\pi$ and $\dfrac{c}{d} \neq e^{a-b}\dfrac{\cos a}{\cos b}$ unless $\cos b = 0$, then $\dfrac{c}{d} \neq e^{a-b}\dfrac{\sin a}{\sin b}$

(c) $b - a = n\pi$ and $\dfrac{c}{d} = e^{a-b}\dfrac{\cos a}{\cos b}$ unless $\cos b = 0$, then $\dfrac{c}{d} = e^{a-b}\dfrac{\sin a}{\sin b}$

EXERCISES 17.2 ■ PAGE 1179

1. $y = c_1e^{3x} + c_2e^{-x} - \frac{7}{65}\cos 2x - \frac{4}{65}\sin 2x$

3. $y = c_1\cos 3x + c_2\sin 3x + \frac{1}{13}e^{-2x}$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

5. $y = e^{2x}(c_1 \cos x + c_2 \sin x) + \frac{1}{10}e^{-x}$

7. $y = \frac{3}{2}\cos x + \frac{11}{2}\sin x + \frac{1}{2}e^x + x^3 - 6x$

9. $y = e^x\left(\frac{1}{2}x^2 - x + 2\right)$

11.



The solutions are all asymptotic to $y_p = \frac{1}{10}\cos x + \frac{3}{10}\sin x$ as $x \to \infty$. Except for y_p, all solutions approach either ∞ or $-\infty$ as $x \to -\infty$.

13. $y_p = (Ax + B)e^x \cos x + (Cx + D)e^x \sin x$

15. $y_p = Axe^x + B\cos x + C\sin x$

17. $y_p = xe^{-x}[(Ax^2 + Bx + C)\cos 3x + (Dx^2 + Ex + F)\sin 3x]$

19. $y = c_1\cos\left(\frac{1}{2}x\right) + c_2\sin\left(\frac{1}{2}x\right) - \frac{1}{3}\cos x$

21. $y = c_1e^x + c_2xe^x + e^{2x}$

23. $y = c_1\sin x + c_2\cos x + \sin x \ln(\sec x + \tan x) - 1$

25. $y = [c_1 + \ln(1 + e^{-x})]e^x + [c_2 - e^{-x} + \ln(1 + e^{-x})]e^{2x}$

27. $y = e^x\left[c_1 + c_2x - \frac{1}{2}\ln(1 + x^2) + x\tan^{-1}x\right]$

EXERCISES 17.3 ■ PAGE 1187

1. $x = 0.35\cos\left(2\sqrt{5}\,t\right)$ **3.** $x = -\frac{1}{5}e^{-6t} + \frac{6}{5}e^{-t}$ **5.** $\frac{49}{12}$ kg

7.



13. $Q(t) = (-e^{-10t}/250)(6\cos 20t + 3\sin 20t) + \frac{3}{125}$, $I(t) = \frac{3}{5}e^{-10t}\sin 20t$

15. $Q(t) = e^{-10t}\left[\frac{3}{250}\cos 20t - \frac{3}{500}\sin 20t\right] - \frac{3}{250}\cos 10t + \frac{3}{125}\sin 10t$

EXERCISES 17.4 ■ PAGE 1192

1. $c_0\sum_{n=0}^{\infty}\frac{x^n}{n!} = c_0e^x$ **3.** $c_0\sum_{n=0}^{\infty}\frac{x^{3n}}{3^n n!} = c_0e^{x^3/3}$

5. $c_0\sum_{n=0}^{\infty}\frac{(-1)^n}{2^n n!}x^{2n} + c_1\sum_{n=0}^{\infty}\frac{(-2)^n n!}{(2n+1)!}x^{2n+1}$

7. $c_0 + c_1\sum_{n=1}^{\infty}\frac{x^n}{n} = c_0 - c_1\ln(1 - x)$ for $|x| < 1$

9. $\sum_{n=0}^{\infty}\frac{x^{2n}}{2^n n!} = e^{x^2/2}$

11. $x + \sum_{n=1}^{\infty}\frac{(-1)^n 2^2 5^2 \cdot \cdots \cdot (3n-1)^2}{(3n+1)!}x^{3n+1}$

CHAPTER 17 REVIEW ■ PAGE 1193

True-False Quiz

1. True **3.** True

Exercises

1. $y = c_1e^{x/2} + c_2e^{-x/2}$

3. $y = c_1\cos\left(\sqrt{3}x\right) + c_2\sin\left(\sqrt{3}x\right)$

5. $y = e^{2x}(c_1\cos x + c_2\sin x + 1)$

7. $y = c_1e^x + c_2xe^x - \frac{1}{2}\cos x - \frac{1}{2}(x + 1)\sin x$

9. $y = c_1e^{3x} + c_2e^{-2x} - \frac{1}{6} - \frac{1}{5}xe^{-2x}$

11. $y = 5 - 2e^{-6(x-1)}$ **13.** $y = (e^{4x} - e^x)/3$

15. No solution

17. $\sum_{n=0}^{\infty}\frac{(-2)^n n!}{(2n+1)!}x^{2n+1}$

19. $Q(t) = -0.02e^{-10t}(\cos 10t + \sin 10t) + 0.03$

21. (c) $2\pi/k \approx 85$ min (d) $\approx$17,600 mi/h

APPENDIXES

EXERCISES G ■ PAGE A12

1. $8 - 4i$ **3.** $13 + 18i$ **5.** $12 - 7i$ **7.** $\frac{11}{13} + \frac{10}{13}i$

9. $\frac{1}{2} - \frac{1}{2}i$ **11.** $-i$ **13.** $5i$ **15.** $12 + 5i$, 13

17. $4i$, 4 **19.** $\pm\frac{3}{2}i$ **21.** $-1 \pm 2i$

23. $-\frac{1}{2} \pm \left(\sqrt{7}/2\right)i$ **25.** $3\sqrt{2}\,[\cos(3\pi/4) + i\sin(3\pi/4)]$

27. $5\left\{\cos\left[\tan^{-1}\left(\frac{4}{3}\right)\right] + i\sin\left[\tan^{-1}\left(\frac{4}{3}\right)\right]\right\}$

29. $4[\cos(\pi/2) + i\sin(\pi/2)]$, $\cos(-\pi/6) + i\sin(-\pi/6)$, $\frac{1}{2}[\cos(-\pi/6) + i\sin(-\pi/6)]$

31. $4\sqrt{2}\,[\cos(7\pi/12) + i\sin(7\pi/12)]$, $\left(2\sqrt{2}\right)[\cos(13\pi/12) + i\sin(13\pi/12)]$, $\frac{1}{4}[\cos(\pi/6) + i\sin(\pi/6)]$

33. -1024 **35.** $-512\sqrt{3} + 512i$

37. ± 1, $\pm i$, $\left(1/\sqrt{2}\right)(\pm 1 \pm i)$ **39.** $\pm\left(\sqrt{3}/2\right) + \frac{1}{2}i$, $-i$





41. i **43.** $\frac{1}{2} + \left(\sqrt{3}/2\right)i$ **45.** $-e^2$

47. $\cos 3\theta = \cos^3\theta - 3\cos\theta\sin^2\theta$, $\sin 3\theta = 3\cos^2\theta\sin\theta - \sin^3\theta$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.